



# Host Map

## *Anoplophora glabripennis*, Asian Longhorned Beetle




**Host Density\***




\* This is a relative risk scale depicting the pest organisms host acreage intensity by county. It is possible to directly compare values between maps of the same type (e.g. Host to Host, Risk to Risk, and NAPPFAST to NAPPFAST).

The U.S. Department of Agriculture's Animal and Plant Health Inspection Service collected the data displayed for internal Agency purposes only. These data may be used by others; however, they must be used for their original intended purposes.

Source: NASS 2007; USFS FIA  
 North America Albers Equal Area Conic (1983)  
 Data contact: Dan Borchert, USDA CPHST, Raleigh  
 Map created: December 2010 by USDA CPHST, Raleigh




1:20,000,000




# Host Map

## *Anoplophora glabripennis*, Asian Longhorned Beetle




# Host Map


## *Anoplophora glabripennis*, Asian Longhorned Beetle




\* This is a relative risk scale depicting the pest organisms host acreage intensity by county. It is possible to directly compare values between maps of the same type (e.g. Host to Host, Risk to Risk, and NAPFFAST to NAPFFAST).



The U.S. Department of Agriculture's Animal and Plant Health Inspection Service collected the data displayed for internal Agency purposes only. These data may be used by others; however, they must be used for their original intended purposes.



1:3,500,000



Source: NASS 2007; USFS FIA  
 North America Albers Equal Area Conic (1983)  
 Data contact: Dan Borchert, USDA CPHST, Raleigh  
 Map created: December 2010 by USDA CPHST, Raleigh