



**Netting to Exclude SWD:  
2013 On-Farm Trials.  
Vern Grubinger**

- **ProTekNet 80 or ProTekNet 25**
- **Highbush blueberries or fall raspberries**
- **'Two cup' red traps with yeast /flour bait**
- **Four locations in Windham County VT**



**bulk batch yeast/flour/sugar bait**



## ProTekNet Options

17 gr    .85 x .85 mm    1-2 yrs

25 gr    .35 x .35 mm    2-3 yrs

60 gr    1.9 x .95 mm    5 yrs

80 gr    1 x .85 mm    7 yrs

Dubois  
Agriculture  
ProtekNet 120gr

Dubois  
Agriculture  
ProtekNet 120gr

Dubois  
Agriculture  
BioTrips 25gr


Dubois  
Agriculture  
ProtekNet Standard 60gr





# Effect of ProTek 80 Netting on SWD Catches in Blueberries

## Grubinger's Blueberry Patch, Dummerston VT












# Effect of ProTekNet 25 on SWD in Raspberries

## Deer Ridge Farm, Guilford VT



# Netting Details

**ProTekNet 80gr** has 1 x .85 mm mesh

- Porosity: 80 % Light Transmission: 83 %
- High Density Polyethylene
- 13' x 328' = \$553 (\$665)

\$5,649/acre @ 7 yrs = **\$807/acre/yr.**

**ProTekNet 25gr** has .35 x .35 mesh

- Porosity: 62 % Light Transmission: 89 %
- Synthetic fiber
- 20' x 820' = \$1,618 (\$1,951)

\$4,298/acre @ 3 yrs = **\$1,433/acre/yr.**

<http://www.duboisag.com/en>



# Effect of Bait Type on SWD catches in Blueberries

## Lost Mile Farm, W. Dummerston VT

