Three Invasive Insects in Vermont, New Hampshire, and Maine


Alan T. Eaton
UNH Cooperative Extension
January 19 - 21, 2016




Brown Marmorated Stinkbug

Discovered in ME 2006 NH 2010 VT 2011









The first BMSB damage in agricultural crops close to New England was apple injury in the southern Hudson Riv. Valley of New York, in 2013.

In 2014, some Connecticut orchards had significant BMSB's and damage.

In 2015 damage was more widespread in Hudson Riv. Valley and CT.







Trissolcus japonicas is a Chinese egg parasite of BMSB being studied by USDA in quarantine, for release in the USA. It was also discovered this year in MD and WA (did not escape). University of New Hampshire Cooperative Extension

SWD (spotted wing drosophila) was discovered in NH/ME/VT Sept. 2011.

NH damage:

2012: \$1,516,000

2013: \$529,000

2014: \$224,000

2015: Similar?



2015 Populations

In both 2014 and 2015, SWD populations in New England increased more slowly than before, and totals were lower than in previous years.


Cooler summer temperatures may have contributed in 2015, and it is possible that natural enemy pressure has increased.


Northern VT/NH/ME has lower numbers than southern VT/NH/ME.

University of New Hampshire Cooperative Extension

Emerald Ash Borer attacks only Ash (Fraxinus). Highly mobile!







Questions?