# Plant-Mediated IPM Systems for Pest Aphids of High Tunnel Vegetable Production



### What's The Big Deal?

### Aphids!!!

- Cause significant crop and revenue loss
  - o Distortion, stunting, sooty mold
- Conventional growers spray chemical pesticides
  - o Threatens human heath and the environment
- Organic growers do nothing or invest in frequent releases natural enemies
  - o Costly



### Project Overview

Develop effective IPM tactics for high tunnel vegetable production suitable for the Northeast and Mid-Atlantic and encourage adoption through a collaborative, multidisciplinary, regional research and outreach initiative ME, PA & VT.

Obj. 1: Develop strategies to sustain aphid nat. enemies in organic high tunnels

Obj. 2: Establish a High-Tunnels Steering Committee

**Obj. 3:** Conduct 3-state (ME, PA & VT) hands-on educational programs

#### Our Cooperators

VT: A. Hazelrigg, Univ. of VT

ME: M. Hutton & J. Lilley, UMaine

PA: E. Sanchez & K. Ayers, Penn State

NY: C. Glenister, IPM Labs





### Research Objective

Test <u>two</u> plant-mediated IPM systems that support biological control of aphids to determine their suitability and effectiveness for <u>two</u> vegetable crop types and <u>two</u> cropping seasons in high tunnels in the Northeast

Cool Season Crops Leafy Greens: Winter




Warm Season Crops
Tomatoes, Peppers, Cucumbers: Summer



### Habitat Plant Systems

#### Summer




Hard Red Spring Wheat Aphid Banker

### Habitat Plant Systems



#### Winter

Dwarf Calendula (*Calendula officinalis* var. Yellow Gem)





Majority Cold Tolerant Max. Height under 18in



Viola (*Viola tricolor* var. Helen Mount) Marigold (*Tagetes patula* var. Little Hero Yellow)

Bush Bean (*Phaseolus vulgaris* var. Provider)

Hard Red Winter Wheat Aphid Banker

### Methods


#### In A Nutshell:

- ✓ Aphid species & numbers
- ✓ Natural enemy types attracted
- ✓ Hyperparasitism
- ✓ Habitat/banker plant performance
- ✓ Costs



# Preliminary Results

Over 700 individual natural enemies encountered Alyssum, Borage, Dill most attractive




### Preliminary Results

Majority were parasitic wasps & mummies, *Orius* adults & nymphs & syrphid adults

Others include various lady beetle life stages, predatory maggots, assassin bugs, lacewing eggs and larvae, etc.

6 species aphids collected from primarily borage



# Summer Habitat Plantings





### Winter Greens







### Questions?



### Thank You!

© 2016 Univ. of VT, Entomology Research Laboratory

This information is based on work supported by the National Institute of Food & Agriculture, US Dept. of Agriculture, Crop Protection & Pest Management Competitive Grants Program, under award #29107. Any opinions, findings, conclusions, or recommendations expressed herein are those of the authors and do not necessarily reflect the view of the US Dept. of Agriculture.