

Quasi-periodicities in the anomalous emission events in pulsars B1859+07 and B0919+06

Haley M. Wahl,¹★ Daniel J. Orfeo,¹ Joanna M. Rankin¹★ and Joel M. Weisberg²★

¹Physics Department, University of Vermont, Burlington, VT 05405, USA

²Physics and Astronomy Department, Carleton College, Northfield, MN 55057, USA

Accepted 2016 June 30. Received 2016 June 30; in original form 2016 April 27

ABSTRACT

A quasi-periodicity has been identified in the strange emission shifts in pulsar B1859+07 and possibly B0919+06. These events, first investigated by Rankin, Rodriguez & Wright in 2006, originally appeared disordered or random, but further mapping as well as Fourier analysis has revealed that they occur on a fairly regular basis of approximately 150 rotation periods in B1859+07 and perhaps some 700 in B0919+06. The events – which we now refer to as ‘swooshes’ – are not the result of any known type of mode-changing, but rather we find that they are a uniquely different effect, produced by some mechanism other than any known pulse-modulation phenomenon. Given that we have yet to find another explanation for the swooshes, we have appealed to a last resort for periodicities in astrophysics: orbital dynamics in a binary system. Such putative ‘companions’ would then have semimajor axes comparable to the light cylinder radius for both pulsars. However, in order to resist tidal disruption, their densities must be at least some 10^5 g cm^{-3} – therefore, white-dwarf cores or something even denser might be indicated.

Key words: radiation mechanisms: non-thermal – pulsars: general – pulsars: individual: B1859+07 and B0919+06.

1 INTRODUCTION

Pulsars are rapidly rotating, highly magnetized neutron stars that emit electromagnetic radiation along their magnetic axes which are tilted with respect to their rotation axes. This circumstance causes their radio signals to sweep across our sightline as the star rotates, producing discrete, regular pulses. Such pulse trains or sequences can be complicated by several known phenomena including *nulls* (one or more consecutive pulses with no detectable power) and *mode changes* (abrupt switches between two distinct emission patterns). Both of these effects have been studied extensively and are broadly consistent with our overall understanding of pulsar action – that is, emission forms cones around the magnetic axis that, in turn, are comprised of a rotating ‘beamlet’ system.

Additional anomalous events were first detected in the emission of radio pulsars B0919+06 and B1859+07 by Rankin, Rodriguez & Wright (2006, hereafter *RRW*). In these events, (1) the regular pulsed emission gradually moved earlier in longitude over a few pulses, (2) remained early for a dozen or more pulses – largely emptying the usual emission window, and (3) then returned over a few pulses to the usual longitude. These events attracted notice because they did not seem to be ‘classical’ mode changes, which exhibit transitions within a single rotation.

RRW examined the events in both pulsars, drawing on six full-polarization Arecibo observations (four of B0919+06 and two of B1859+07 made between 1981 and 2005). They attempted to work out the two pulsars’ emission geometry and examined several possible explanations for the events. The possibility of some new type of mode changing was examined, but was ruled out because of the gradual onsets and relaxations of the events. Changes in emission altitude were also examined, but the needed displacements within the magnetosphere seemed unreasonably large. Appeals to ‘absorption’ or other effects seemed ad hoc. In short, *RRW* were unable to identify a plausible physical cause for these observed events. However, they did show that both the normal and event profiles seemed to stem from partial illumination of the stars’ polar caps, as artificial profiles constructed from a comparable number of normal and event pulses produced core-cone triple profiles with the usual core and conal angular dimensions (Rankin 1993a,b).

Several other studies have drawn attention to these two pulsars. Mitra & Rankin (2011) conducted sensitive single pulse polarimetry on most of the pulsars originally categorized by Lyne & Manchester (1988) as ‘partial cones’ – including B0919+06 – and their analysis tends to support the interpretation of *RRW* to the effect that the events entail a partial illumination of the pulsar’s emission cone, late for the normal emission and earlier during the events. Evidence for long term periodicities on the order of tens of years in the timing of B0919+06 were developed by Lyne et al. (2010) and Shabanova (2010) and thoroughly studied recently by Perera et al. (2015). A

* E-mail: haley.wahl@uvm.edu (HMW); Joanna.Rankin@uvm.edu (JMR); jweisber@carleton.edu (JMW)

similar study on B1859+07, again drawing on the unexcelled pulsar timing archive at the Jodrell Bank Observatory, has been carried out by Perera et al. (2016). Most recently, Han et al. (2016) published the results of ~ 30 h of B0919+06 observations using the Jiamusi 66-m Telescope and found events similar to those identified by RRW.

In this paper, we undertake a new study of these emission events in pulsars B0919+06 and B1859+07. In order to emphasize their distinct character, we refer to the events as ‘swooshes,’ returning to the (unpublished) terminology used by RRW in the preparation of their paper. First noticed in the 1990s in Arecibo observations of B0919+06, the phenomenon was ignored until it was again seen by RRW – and now in this work with three different Arecibo instruments. That the events were also seen with the Jiamusi Telescope rules out any instrumental cause. After initial analysis of these two pulsars by RRW, it had become clear that an entirely new approach was needed to investigate and explain the swoosh phenomenon, and we began by carrying out much longer and more extensive Arecibo observations of both pulsars. Our analyses produced evidence of a perplexing periodicity in both stars, which in turn opened up new potential avenues of interpretation. Section 2 describes the observations. Sections 3 and 4 examine events in each star in more detail, and Sections 5 and 6 discuss the evidence for their periodicity. Sections 7, 8, and 9 then discuss possible mechanisms for the swooshes. Finally, Section 10 summarizes the results and discussion.

2 OBSERVATIONS

These new observations were carried out using the upgraded Arecibo Telescope in Puerto Rico with its Gregorian feed system, 327-MHz (*P* band) or 1100–1700-MHz (*L* band) receivers, with either Wideband Arecibo Pulsar Processors or Mock spectrometer backends. At *P* band, four 12.5-MHz bands were used across the 50 MHz available. Four nominally 100-MHz bands centred at 1170, 1420, 1520, and 1620 MHz were used at *L* band, and the lower three were usually free enough of radio frequency interference such that they could be added together to give about 300-MHz bandwidth nominally at 1400 MHz. The four Stokes parameters were calibrated from the auto- and cross-voltage correlations computed by the spectrometers, corrected for interstellar Faraday rotation, various instrumental polarization effects, and dispersion. The resolutions of the observations are all about a milliperiod. The Gregorian and modified Julian dates as well as the lengths of the observations are listed in Table 1.

3 CHARACTERISTICS OF THE SWOOSH EVENTS

The primary signature of the swooshes in pulsars B0919+06 and B1859+07 is a shift – to an earlier longitude – at which the emission occurs. As seen in Fig. 1, the shifts during these swooshes appear nearly identical in B0919+06 and B1859+07 when examined on an interval of 1000 or more pulses. However, some differences are perceptible between the events in each pulsar when they are examined in closer detail.

Fig. 1 exhibits key features of the swooshes in the two pulsars: one can see that the swoosh transitions are always in the earlier direction and that both onset and termination are gradual over a few pulses. Also, the polarized character of the emission remains very similar during a swoosh. L/I and V/I show no systematic change, and χ rotates linearly even while the peak emission moves to earlier longitudes. Finally, there is generally little or no change in

Table 1. Observational parameters.

Pulsar RF (MHz)	Date (yr/m/d)	MJD	Length (pulses)	Number of events
B0919+06				
1425	2003/8/03	52854	1115	2
327	2003/10/04	52916	4180	1
325.85	2013/5/05	56417	17 259	6
325.85	2013/10/11	56576	17 810	7
1392.5	2013/10/12	56577	13 994	4
1494	2015/8/23	57257	16 558	6
2250	2015/4/18	57130	2088	1
2250	2015/7/12	57215	53 053	22
2250	2015/7/14	57217	67 611	21
2250	2015/8/17	57251	65 003	26
2250	2015/8/18	57252	35 306	16
2250	2015/8/19	57253	25 001	6
B1859+07				
1525	2003/04/10	52739	1021	~ 7
1520	2005/1/2	53372	2096	~ 12
1392.5	2013/3/26	56377	6460	~ 40
1392.5	2015/4/8	57121	12 195	~ 171
1394	2015/12/5	57361	12 154	~ 141

the intensity of the individual pulses during a swoosh, but missing pulses are seen in B0919+06, see details below, which is why we choose to call these events ‘swooshes’ rather than ‘flares.’

3.1 B0919+06

Our analysis of B0919+06 focuses on 12 observations detailed in Table 1. Half of the pulse sequences are taken from our own Arecibo observations while the other half was acquired by Han et al. (2016). Using the 1000-pulse displays, (see Fig. 1) these events stand out as -10° shifts in the longitude of the emission window that persist for anywhere from 32–70 pulses. Though the length and extent of the shifts can vary, B0919+06 events follow a consistent pattern: first, weaker emission is seen for 2–5 pulses which can sometimes be so drastic as to appear as a null. Secondly, during the maximum longitude shift of the next several dozen pulses, the usual emission region largely empties, the polarization angle exhibits a modest progressive rotation of up to about -18° , and L/I decreases from some 80 per cent to 50 per cent. Finally, the emission gradually returns to its usual longitude over a few pulses, sometimes followed by another weakening or apparent nulling for 2–5 pulses. This pattern is consistent for each B0919+06 event throughout all of our observations, following the same pattern over the more than three decades of Arecibo observations at all frequencies. Though the signatures of each event are almost identical on a gross scale, Han et al. were able to classify them into four subtly different categories (see Section 4 for details).

3.2 B1859+07

The B1859+07 events appear almost identical to those of B0919+06 when viewed on the 1000-pulse display of Fig. 1. However, upon closer examination, the B1859+07 swooshes are seen to have much more varied and irregular patterns: their duration can be anywhere from 20 to 120 pulses and they appear to have different morphological characteristics (see Section 4). They are also not always fully distinct and separated from each other, which can make it difficult to identify individual events. This circumstance is the




Figure 1. Single-pulse polarization displays of the swoosh events in 1000-pulse intervals for both pulsars: two swooshes can be seen in B0919+06 (left) and some seven in B1859+07 (right). The total power I , fractional linear L/I , polarization position angle χ , and fractional circular polarization V/I are colour-coded in each of four columns according to their respective scales at the left of the diagram.

origin of the uncertainty in the number of B1859+07 events listed in Table 1. These intervals of ‘connected’ events appear a few times in the early shorter observations taken over the course of about 10 yr but increase in frequency in the later MJD 56377 observation. The even-more-recent 2015 April observation (MJD 57121) shows a further increase in the frequency of these irregular, consecutive events, and events in general, to the point where the number of swooshes in each 1000-pulse interval almost doubled compared to relatively consistent tallies in the previous observations. A long 2015 December observation also showed an increase in the frequency of these events compared to observations from the previous decade, though the event frequency is slightly below that of the 2015 April observation. Further investigation with longer observations is needed to examine the cause of this sudden increase in event frequency.

4 DO EACH OF THE SWOOSHES LOOK THE SAME?

In *RRW*, all swooshes studied in both B0919+06 and B1859+07 seemed to exhibit one specific pattern in which the emission gradually decreased in longitude, stayed in that position for a number of pulses, and made a gradual transition back to the normal emission longitude. *RRW* concluded that the events in both stars followed this pattern, but upon further analysis on longer observations, it was discovered that the events in B0919+06 as well as B1859+07 can come in many different shapes and sizes. Han et al. were able to categorize the events in B0919+06 into four categories: Π Type (where the emission abruptly jumps to an earlier longitude, stays

there for a few tens of periods, and then abruptly jumps back down to its normal longitude), M Type (where the emission moves to an earlier longitude for some 20 periods and then gradually returns back) (see Fig. 2 for an example), Δ Type (where the emission sharply decreases in longitude and then sharply transitions back without remaining in the early state for more than 1–5 pulses), and λ Type (where the emission moves sharply to an earlier longitude and then slowly transitions back to the regular state).

Just as Han et al. were able to categorize the shapes of events seen in B0919+06, we were able to perform similar analyses on the B1859+07 swooshes and sort them into seven categories based on their morphology. Due to the fact that the B1859+07 swooshes are very similar to those of B0919+06 when analysed in detail, we take from the Han et al. analysis the first four morphological categories of B1859+07 swooshes: Π , M , Δ , and λ Types. The last three B1859+07 swoosh categories are patterns which do not appear in any current B0919+06 observations: two brief consecutive shifts to an earlier longitude which essentially appear as two connected Δ Type events with no more than 1–5 pulses between them (where the emission may or may not return to its original longitude in the middle), we refer to as ω Type; events which feature brief sharp transitions to an earlier longitude like a Δ Type except fill the window from starting longitude to ending longitude during the event, we refer to as Δ Type; and finally the O Type, which are similar to the Δ Type but have slower transitions causing the entire event to appear rounded.

Fig. 3 gives a 200-pulse display showing two of these events in MJD 56377: an ω Type and an M Type event. Of additional note in




Figure 2. Higher resolution 200-pulse display showing the second B0919+06 swoosh in Fig. 1.




Figure 3. Higher resolution 200-pulse display of the fifth and sixth major B1859+07 swooshes in Fig. 1.

this observation are several long intervals in which no events occur, long stretches of ω Type events, single-pulse nulls, and longer nulls lasting 2–10 pulses. Analysis of the shorter observations (MJD 52738 and 53372) reveals additional irregular behaviour. The fact that the categorizations in both our analysis and the Han et al. analysis seem very similar and the types of events look very much the same when studied closely at small scales strongly indicates

that a similar phenomenon is causing the events in both pulsars and therefore allows them to be studied side by side.

5 IDENTIFICATION OF A COMPLEX SWOOSH PERIODICITY IN PULSAR B1859+07

Though the events in B0919+06 and B1859+07 appear to be somewhat different when studied in detail, they both exhibit a similar gradual early shift pattern of the emission longitude. As seen in Table 1, the events seem to appear much more frequently in B1859+07 than they do in B0919+06, which prompted further analysis of the former star. Moreover, the B1859+07 swooshes give the impression of occurring with a rough regularity of several hundred pulses as we see in Fig. 1.

We therefore first studied the MJD 56377 observation of B1859+07 by computing a longitude-resolved fluctuation spectrum, so as to investigate the reality of any event periodicity. This spectrum, shown in the top panel of Fig. 4(a), remarkably displays a well-defined periodic feature with a peak frequency at 0.00685 cycles period⁻¹ ($P = 146$ pulsar rotation periods) associated most strongly with the leading and trailing edges of the profile. The latter property ties this principal fluctuation to the events, since a longitudinal shift of emission would be most evident at these longitudes. A more accurate value for the total fluctuation frequency can be computed by using all of the spectral components with intensities greater than the half-power point (Δf is this half-power width), giving $P = 153 \pm 3$ periods. Note, however, that the ‘ Q ’ [$=f/\Delta f$] of the feature is not very large, perhaps 2, so the effective error is much larger than that given just above. This squares with our further efforts to measure the period of the swooshes with a periodogram analysis, which showed little regularity in the intervals between them.

Indeed, it is difficult to understand how the strong feature in the MJD 56377 fluctuation spectrum arises when examining the distribution of intervals between swooshes. The majority of the intervals fall between 50 and 100 periods. However, there are several long intervals without swooshes, one 280 and another 351 periods.

None the less, given the identification of this fairly regular modulation in our MJD 56377 observation, it seemed important to see what our other observations showed. The total fluctuation spectra of the MJD 52739, 53372, 57121, and 57361 observations are shown in Fig. 4(b)–(e), respectively; and given their shorter length or more complex modulation, fast Fourier Transforms of length 512 were used for them, while a length of 1024 was employed for the results of Fig. 4(a). The short MJD 52739 observation of Fig. 4(b) shows a strong feature at 0.0068 cycles period⁻¹ with a period of 148 ± 5 stellar rotations. There is also power at 0.014 cycles period⁻¹ or 73 ± 3 periods, which seems to be a harmonic of the main feature. The somewhat longer MJD 53372 spectrum of Fig. 4(c) has two strong peaks at 170 and 102 periods; however, when all of the fluctuation power in the first seven components is weighted and averaged, the result is 152 ± 8 periods, again with a small Q . This all said, we re-emphasize that the feature widths are all much larger than the errors in the spectral peaks given just above, or equivalently their ‘ Q ’s are again of the order of 2. Therefore, it seems that the swooshes have a consistent average period while exhibiting substantial irregularities in their intervals of occurrence.

The total fluctuation spectra of the MJD 57121 and 57361 observations (Figs. 4d–e) show a somewhat different spectrum. This was not too surprising given the above discussion of their more frequent and difficult to interpret swooshes. Both are also the longest




Figure 4. Fluctuation spectra of B1859+07, showing evidence for event periodicity. (a) The display for MJD 56377 contains three contiguous panels, with the pulse profile on the left, the longitude-resolved fluctuation spectrum in the top right, and the overall fluctuation spectrum at the bottom; while the overall fluctuation spectra for MJD 52739, 53372, 57121, and 57361 are shown in (b)–(e), respectively. The profile and longitude-resolved spectra are shown in the left-hand panels and the total fluctuation spectrum in the lower one. Note that all of the overall fluctuation spectra are plotted only to 0.15 cycles period⁻¹. The feature at 0.006 85 cycles period⁻¹ corresponds to a periodicity of 146 rotation periods. Fourier transforms of length 1024 were used for the MJD 56377 analysis and 512 for the others. The different amplitude scales result from use of the two different backend processors.

of our observations at more than 12 000 pulses. Fluctuation power in these observations is spread over a wider range than in the other observations. Both spectra show two peaks. In the 57121 spectrum, one is at 170 and the other at 85 periods; whereas the 57361 has peaks at 143 and 81 periods, all with errors of some 10 periods. A weighted average of the 57121 spectral power up to the sharp decrease at about 0.014 cycles period⁻¹, gives a period of some 144 ± 10 rotation periods. For the 57361 spectrum, the errors are such that the two features could be harmonically related as indicated above, though all the Qs are small.

6 ARE THE SWOOSHES IN B0919+06 QUASI-PERIODIC AS WELL?

The swooshes in B0919+06 are much less frequent than they are in B1859+07. In our six observations, totalling some 8.5 h and 70 916 pulses, we have identified only 26 swoosh events, while Han et al. were able to identify a total of 92 abnormal emission events in some 30 h or 248 062 pulses total.

The three shortest intervals between swooshes in the full set of B0919+06 observations are 292, 495, and 549 pulses. The next largest group are 728, 779, 806, and 880. All the others are much longer, the 4180-pulse MJD 52916 sequence shows but a single swoosh, and in two instances, the interval is longer than 10 000 pulses. Previous observations taken with the Arecibo Telescope showed shortest intervals between swooshes as 662, 837, and 900 pulses with the next largest group being 1508, 1742, 1924, 1529, and 1341 pulses.

A rough approximate period using just our Arecibo observations was calculated to be about 700 pulses or 300 s. With the introduction of the Han et al. sequences, the B0919+06 events could be studied over much longer time-scales, which revealed that their occurrence is at least as complex as in B1859+07. For B1859+07, a consistent long-term periodicity of some 150 periods is found using Fourier techniques, whereas mapping of the intervals between events shows many smaller and larger intervals. We have investigated the Han et al. B0919+06 sequences using Fourier methods as well, and we find no indication of periodicity at all, only low-frequency fluctuation power at periods greater than several hundred stellar rotations.

7 HOW CAN WE INTERPRET THE SWOOSH PHENOMENON?

The putative periodicity in the swoosh events in pulsar B1859+07, introduces a variety of new directions and complications into any effort to interpret the swooshes physically. Let us summarize –

(i) the swooshes are not conventional mode changes for a variety of reasons. They typically begin and end gradually. No other pulsar switches modes by simply altering the longitude range of its emission (see Rankin 1986 for references to studies of the then known mode switching pulsars).

(ii) the swooshes always represent intervals of earlier emission. RRW’s suggestion of ‘partial conal’ emission remains possible, but we have no physics to understand why the illumination would usually be late but then earlier during the swooshes (See Mitra & Rankin 2011 for a full study of ‘partial cone’ emission).

(iii) that the swooshes are caused by an emission-height change remains possible, but we think unlikely. In no pulsar has this effect been clearly demonstrated or a physical cause identified. Ad hoc assumptions are needed to estimate such a change and again, why such a height change should occur (e.g. why the height should increase rather than decrease).

(iv) no emission mechanisms are known that would produce quasi-periodic shifts of the sort observed in the swooshes. Some other emission effects are known with very long time-scales over multihour long time-scales (e.g. the carousel rotation slowdown after B-mode onset in pulsar B0943+10; see Rankin & Suleymanova 2006), but no periodicity has been identified with effects such as these.

(v) pulsations, a form of surface oscillations known to exist in white dwarfs, have been explored as a possible cause for periodic phenomena in pulsars (e.g. Rosen & Clemens 2008). In terms of

the events in our pulsars, we do not believe that oscillations are the cause for three reasons: (1) there is no conclusive evidence that these oscillations/pulsations exist in pulsars; (2) the expected \sim ms oscillation time-scales would be much smaller than the \gtrsim 1 s time-scales of our events; and (3) such oscillations tend not to have a constant period but quicken as they lose energy, which is not seen in our swooshes.

8 COULD THE SWOOSHES REFLECT AN ORBITING BODY OR BODIES?

Let us consider an alternative explanation. In astrophysics, a fundamental physical cause of periodicity is orbital dynamics. However,

(i) periodicities due to orbital causes tend to have a high Q so it is difficult to understand how a ‘companion’ orbiting the pulsar could cause the very rough periodicities we apparently see in B1859+07 and B0919+06.

(ii) the time displacements are not easy to understand: the swooshes move the emission earlier in both pulsars. A highly eccentric orbit might produce a displacement in one direction for one part of the orbital period, but it is likely that secular variations of the orbital elements would eventually produce shifts in both directions.

(iii) binary orbits are not known for such short periods: 153 rotation periods for B1859+07 would be only 99 s, and 700 rotations for B0919+06 only 300 s.

These conclusions, however, strictly follow only if the orbit is a simple binary. More complex orbital effects follow from more complex systems, including the possibility of a chaotic system. Such close orbital mechanics would certainly entail decay by gravitational radiation, so the finite lifetime of the system must be considered.

9 WHAT IF THE SWOOSHES DID REFLECT AN ORBITING BODY OR BODIES?

Given that we cannot explain the swooshes and their putative periodicities by other means, we here explore what kind of orbital systems might produce the effects we observe. Specifically, we describe the consequences of interpreting the inter-swoosh quasi-period P_s as the orbital period P_b of such systems.

If the companion had a mass much smaller than that of the neutron star, we could use Kepler’s Third Law $(M/M_\odot)(P \text{ yr}^{-1})^2 = (a_c/\text{AU})^3$ to compute the companion’s orbital semimajor axis a_c , given a system mass $M \sim 1.4M_\odot$ and orbital period P . For PSR B1859+07, the orbital period $P = P_s$ is ~ 150 s or 2.1 μy , so that $a_c \sim 180 \mu\text{AU}$ or some 27 000 km. Remarkably, this is only a little larger than the pulsar’s light-cylinder radius of 20 560 km! If we do a similar calculation for B0919+06 using the period of about 700 pulses, we find an a_c value of 760 μAU or some 120 000 km, as compared to a light cylinder of radius 30 560 km.

These estimates show that the semimajor axes of the orbits are larger, but only a little larger, than the light-cylinder radius for both pulsars. Were the orbits elliptical, the ‘satellite’ could pass close to or even inside the light cylinder. It is also interesting that the closer orbit to its light cylinder is that of B1859+07, where we apparently see more of the swooshes than for B0919+06. Given that the orbital period of the ‘satellite’ and the rotation period of the pulsar are asynchronous, one can further speculate that a variety of effects might be produced, as a function of the (unknown) orientation of the ‘satellite’ with respect to our sightline when it passed close to the pulsar.

The lack of cyclical emission-longitude variations having periodicity P_s (aside from the low duty-cycle swooshes themselves) indeed indicates that the companion mass must be relatively low, thereby validating the assumptions underlying the above calculations. Even if the swoosh longitude offsets themselves are interpreted as the pulsar reflex motion, their ~ 10 ms amplitudes indicate a projected pulsar semimajor axis $a_p \sin i$ of only ~ 10 light-ms, which is much less than the characteristic size of the companion orbit, as calculated above.

Could any companion persist in such an environment so close to the pulsar? Tidal disruption would be a major issue, and the ability of a ‘satellite’ to resist disruption provides a constraint on its density. Disruption occurs when $M/a_c^3 > \rho_c$, where ρ_c is the companion’s density (Harwit 1973). For B1859+07, this limit on the density is about 10^5 g cm^{-3} , suggesting that only a body having a density of a white dwarf core or greater could so persist. Indeed, pulsar-white dwarf binaries are known, but with much longer periods; however, one such system with a period of 2 h has been identified (Bailes et al. 2011), while white dwarf-white dwarf binaries with orbital periods as short as 12.75 min have been discovered (Brown et al. 2011). Furthermore, the lack of observed binary systems of shorter known periods could easily be due to the difficulty of detecting them.

10 SUMMARY

We report extensive new Arecibo observations and analyses studying the strange swooshes in pulsars B0919+06 and B1859+07. We find much more complexity than was earlier reported by Rankin et al. (2006); however, we confirm the basic effect involving usually gradual, temporary, about 10° shifts in the emission window to earlier longitudes. Here also, we provide some evidence for a quasi-periodicity in pulsar B1859+07 and perhaps in B0919+06. We examine a number of effects that might produce the swooshes and find no obvious mechanism. In particular, the effects might represent mode changes, but no example of a gradual mode change is known, so swooshes appear to be some other phenomenon. These difficulties have led us to explore whether some type of co-orbiting companion with an unprecedentedly short period and separation might be involved. Such a system would entail so many effects of tide and heating that it is difficult to predict what the consequences might be. Given that WD–NS and WD–WD binaries are known with periods 10–100 times larger, one can ask what happens as those systems evolve.

ACKNOWLEDGEMENTS

HMW gratefully acknowledges a Sikora Summer Research Fellowship. Much of the work was made possible by support from the US National Science Foundation grant 09-68296 and from a Vermont Space Grant. JMW acknowledges support from NSF Grant AST-1312843. We thank Han et al. 2016 for access to their total-power data on PSR B0919+06. The Arecibo Observatory is operated by SRI International under a cooperative agreement with the National Science Foundation (AST-1100968), and in alliance with Ana G. Mendez-Universidad Metropolitana, and the Universities Space Research Association.

REFERENCES

- Bailes M. et al., 2011, *Science*, 333, 1717
Brown W. R., Kilic M., Hermes J. J., Allende Prieto C., Kenyon S. J., Winget D. E., 2011, *ApJ*, 737, L23

- Han J. et al., 2016, MNRAS, 456, 3413
Harwit M., 1973, *Astrophysical Concepts*. Wiley, New York
Lyne A. G., Manchester R. N., 1988, MNRAS, 234, 477
Lyne A. G., Hobbs G., Kramer M., Stairs I., Stappers B., 2010, *Science*, 329, 408
Mitra D., Rankin J. M., 2011, ApJ, 727, 92
Perera B. B. P., Stappers B. W., Weltevrede P., Lyne A. G., Bassa C. G., 2015, MNRAS, 446, 1380
Perera B. B. P., Stappers B. W., Weltevrede P., Lyne A. G., Rankin J. M., 2016, MNRAS, 455, 1071
Rankin J. M., 1986, ApJ, 301, 901
Rankin J. M., 1993a, ApJ, 405, 285
Rankin J. M., 1993b, ApJS, 85, 145
Rankin J. M., Suleymanova S. A., 2006, A&A, 453, 679
Rankin J. M., Rodriguez C., Wright G. A. E., 2006, MNRAS, 370, 673 (RRW)
Rosen R., Clemens J. C., 2008, ApJ, 680, 671
Shabanova T. V., 2010, ApJ, 721, 251

This paper has been typeset from a \TeX/L\TeX file prepared by the author.