



A Report from the University of Vermont Transportation Research Center

Spatial Extent of the Impact of Transported Road Materials on the Ecological Function of Forested Landscapes

TRC Report #12-012 | Neher, Asmussen and Williams | August 2012

Spatial Extent of the Impact of Transported Road Materials on the Ecological Function of Forested Landscapes

UVM Transportation Research Center

August 17, 2012

Prepared by:

Deborah A. Neher, Ph.D.
David Asmussen, M.S.
Kristin Williams, M.S.

Department of Plant & Soil Science
Jeffords Hall
63 Carrigan Drive
Burlington, VT 05405

Phone: (802) 656-2630
Website: www.uvm.edu/~dneher

1. Acknowledgements

We thank Sarah T. Lovell for the initial project idea and the USDOT DTRT06-G-0018 for funding this work through the University Transportation Center (UTC) at the University of Vermont. We also thank Jim Cota at Vermont Agency of Transportation, Jim Sullivan at the University of Vermont Transportation Research Center, Hayden Lake at Cohosh Forestry, Elizabeth Thompson, Cathy Paris, Josef Görres, Alan Howard, Thomas R. Weicht, Donald S. Ross, Kimberly Wallin, and Nicolas Le Blanc for consultation and technical assistance.

Disclaimer

The contents of this report reflect the views of the authors, who are responsible for the facts and the accuracy of the data presented herein. The contents do not necessarily reflect the official view or policies of the UVM Transportation Research Center. This report does not constitute a standard, specification, or regulation.

Table of Contents

Acknowledgements and Disclaimer
List of Tables and Figures

1. Introduction	5
2. Research Methodology.....	6
2.1 Site Selection.....	8
2.2 Data collection.....	9
2.2.1 Soil Chemistry.....	9
2.2.2 Soil Biology	10
2.2.3 Vegetation	11
2.2.4. Road Attributes	11
2.3 Analytical Methodology	11
3. Results	12
4. Implementation/Tech Transfer	19
5. Conclusions.....	20
6. References.....	22
7. Appendices	25
8. Equations.....	32

List of Tables

Table 3-1. Soil properties for each of three road types page 15
Table 3-2. Mean values of soil component from all road types page 16

List of Figures

Figure 2-1. Map of watersheds.....page 7
Figure 2-2. Transect layout.page 8
Figure 2-3. Measured Dimensions of Roadside Microtopographypage 9
Figure 3-1. Mean levels of lead (mg/kg) in roadside soil based upon categorical distance from road..... page 13
Figure 3-2. Mean levels of Calcium (mg/kg) in roadside soil based upon categorical distance from road..... page 14
Figure 3-3. Mean levels of Sodium (mg/kg) in roadside soil based upon categorical distance from road..... page 17
Figure 3-4. Mean MI index in roadside soil based upon categorical distance from road page 18
Figure 3-5. Mean ratio of fungal feeding to fungal and bacterial feeding nematodes in roadside soil based upon categorical distance from road..... page 19
Figure 4-1. Schema of projects integrated together under TRC Signature 1. page 20

1. Introduction

Roads have varied ecological impacts on the adjacent plant and soil environment due to physical and chemical disturbances resulting from roadway construction, roadside maintenance, and vehicle deposition. The two main areas influenced by a road are the roadside right-of-way and vegetated region just beyond the right-of-way, which often consists of a semi-natural habitat with some native species [1].

Roadway construction is a major disturbance that has direct and indirect impacts on plant communities and soil properties. The initial clearing for the road corridor during the construction phase typically establishes the base age of woody species [2]. Nearby vegetation has a large influence on the species richness of the vegetation that repopulates the cleared areas [3,4,5]. Site grading during the clearing process alters the hydrology of the roadside environment, resulting in channelization of streams, draining of wetlands, or development of new hydrologic zones which can create or destroy habitat for various plants [6,7,8]. Finally, as the road base is built up, large quantities of material are imported, creating a source of mineral material unique to the local environment that may contribute to later chemical disturbances [9].

In addition to the initial physical disturbance from the construction of the road, vehicular traffic and regular maintenance of the road and right-of-way causes recurring physical disturbance. Regular annual mowing will select for plants that seed earlier in the season (prior to the mowing date) and plants with a lower growth habit [10]. Vehicle-generated wind currents can act as dispersal agents for certain species of plants [5,11], sometimes spreading invasive species [8,12]. Mobilized dust in wind currents can travel for hundreds of meters, settle on plants, and interfere with photosynthesis and transpiration [9], which could select for species more tolerant or adapted to this roadside situation.

The roadside environment is also influenced by chemical disturbances, which primarily impact the soil, and indirectly impact vegetation. Though most reports investigate chemical disturbance from recurring processes such as dust deposition, road salt, and exhaust [9,13], roads also have initial chemical impacts due to leaching of new construction material. The first flush of chemicals from a new road may contribute to the current levels of chemicals in roadside soils if the particular compound is retained in the soil [14]. Acute chemical disturbance can also occur after a road is constructed through accidental spills of hazardous materials or passenger vehicle accidents may leak materials [15].

Road dust and seasonal salt applications are important contributors to recurring chemical disturbances that continuously impact plant communities

and soil health. Road dust is composed of finely ground minerals of the road's parent material. If a road base is calcareous (particularly with gravel roads), the resulting dust can significantly change the pH next to the road that, in turn, alters the availability of micronutrients [9]. On asphalt roads, the dust can contain ground particles of tires, brake lining, and asphalt. When distributed to the roadside environment, these materials can contribute to higher levels of heavy metals, particularly zinc [16]. In areas with snow, highway departments apply road salt seasonally to reduce ice and allow faster and safer movement of vehicles. This salt is transported to roadside soil during winter when large particles are knocked off the road by vehicles, or in spring thaws when salt goes into solution during transport with melted snow [7]. Salt in the roadside environment can cause water stress in plants and, if washed from the soil by precipitation, it will travel great distances through surface aquatic systems and potentially to large bodies of water [6].

While there have been studies investigating different aspects of the roadside environment, there is a need for research in forest ecosystems and for development of methods to predict roadside environmental conditions with distance and road use intensity. This study determines how roads within a northern hardwood forest change the native plant and soil conditions at various distances from the road. It also provides a method to predict plant and soil conditions based on traffic volume. The study makes an important contribution to our understanding of plant-soil interactions, as influenced by roadside conditions. Specifically, the objective of this study was to determine the spatial extent of the effects of the road and these transported materials on forest plant communities, soil chemistry, and soil nematode communities immediately surrounding roadways broadly classified as 'highways,' 'two-lane paved,' and 'gravel' which correspond roughly with the Federal Highway Administration's classification of arterial, collector and local [17]. We propose to use nematodes as bioindicators because of their ubiquity, known response to chemical and physical perturbations to soil and water, and current consideration in regional and national monitoring programs. They integrate chemical and physical properties and the microbial community at lower positions in the food chain [18].

2. Research Methodology

The study was conducted in Chittenden County, Vermont, in the Northeastern U.S., where deicing salts are spread regularly on roads during winter months. Land cover in the state of Vermont is dominated by forest (approximately 75%), and many of the forested areas are directly connected with the roadways. The forests within study areas are generally classified as Northern Hardwood Forests and White Pine Northern Hardwood Forests [19, 20]. Sample sites were located throughout Muddy Brook (8,262 ha) and Allen Brook (2,900 ha) watersheds (Fig. 2-1) in mostly rural residential areas [21].

These locations were chosen to co-locate our study with a concurrent study by a team of aquatic ecologists investigating water quality, stream integrity and water pollutant load in relation to road type and road traffic density. The long-term goal was to link the watershed results with roadside results.




Figure 2-1. Map of watersheds and sampling locations for this study. Photo Credit D. Asmussen.




Figure 2-2. Transect layout.

2.1 Site Selection

Potential sampling sites were determined using aerial photography in Google™ Earth version 6.03.2197 to locate sections of forest that covered an area at least 100 m perpendicular to the road, and extending at least 200 m parallel to the road (Fig. 2-2). The distance was chosen to insure independence among transects by avoid overlap or interference. Ten locations were chosen for each road type from the potential sample sites using a random sequence generator ^[22] to minimize bias. These GPS points recorded by the ground survey were loaded into ESRI ArcMap program for analysis with other map layers.

The sampling strategy within a transect was adapted from a study of roadside vegetation in Terra Nova National Park that defined modified roadway zones as the shoulder, sideslope, ditch, backslope, and native vegetation ^[23]. The shoulder is located next to the driving surface of the road and the sideslope is the adjacent area built up during road construction to support the main road surface. The ditch is a low point that carries away water from the road surface and the backslope is the cleared area that maintains ditch functions.




Figure 2-3. Measured Dimensions of Roadside Microtopography. Tree icon represents the edge of the forest which was defined as visually in line, and parallel the road, with the first tree > 8 cm diameter at breast height (DBH). The road crown (0m) is effectively the center of each road type.

2.2 Data Collection

Soil samples were collected from the three road types, at the center of the six microtopographic locations from 11 May to 3 June 2009. Samples were taken at six categorical distances based upon roadside micro-topography along each transect: a) the road shoulder edge; b) the middle of the sideslope; c) the ditch trough; d) the middle of the backslope; e) 10 meters into the forest from the forest edge; f) and 50 meters from road crown (Fig. 2-3). In sum, 216 samples (3 road types x 12 transects x 6 distances) were taken to analyze chemical and physical soil properties. In addition, soil nematode communities were taken along 10 of the 12 transects totaling 180 samples. In the forest, leaf duff and large woody debris were gently removed to reveal the organic horizon (O₂) before sampling.

2.2.1 Soil Chemistry. At each sampling location, two soil cores were collected using an intact soil corer (5 cm diameter, 7.6 cm length) at 1 m on either side of the predetermined location parallel to the road. Each of these soil cores were placed in a plastic bag, labeled and stored at 4 °C until they could be

processed. The samples were air-dried and passed through a 2-mm sieve prior to analysis.

Soil nutrients were extracted for analysis using a Modified Morgan method [24, 25]. Roadside soils were analyzed for nutrients, including available phosphorus (P), potassium (K), magnesium (Mg), aluminum (Al), calcium (Ca), zinc (Zn), sulfur (S), manganese (Mn), boron (B), copper (Cu), iron (Fe), sodium (Na), lead (Pb), nickel (Ni), cadmium (Cd), chromium (Cr), cation exchange capacity (CEC) and cation ratios of calcium, potassium and magnesium (% Ca, % K, % Mg). The pH was determined using a Mehlich buffer method with water. CEC was calculated from the % Ca, % K and % Mg. The percent organic matter (% OM) was determined by loss on ignition and converted to a Walkley-Black equivalence [26]. Chemical analysis was performed at the University of Vermont Agricultural and Environmental Testing Lab (http://www.uvm.edu/pss/ag_testing/).

2.2.2 Soil Biology. Nematodes were collected using a Dutch soil auger (5-cm diameter) to the depth of 30-cm, or as deep as possible in the case of compaction or gravel. An Oakfield probe was highly ineffective with roadside soils due to compaction and the high content of large rocks and gravel. Each nematode sample was a composite of five soil cores taken approximately one meter apart (one pace), in a line parallel to the road and perpendicular to the transect (one on the transect and two in either direction from the transect). Given the roadside conditions, some shoulder samples required more than five cores to obtain an equivalent volume of soil to reduce any bias associated with sampling effort or volume.

Soil nematode samples were mixed in a bucket, and approximately 600 g were then placed in plastic resealable bags and stored in an insulated container to maintain field temperatures and moisture. In the laboratory, soil samples were stored at 16° C in a mechanical convection incubator until soil extraction was completed.

Nematodes were extracted from 200 cm³ of each soil sample using a modified Cobb's decanting and sieving method [27], followed by cotton-wool filter extraction method tray [28] for 48 hrs. To extract nematodes, the soil sample was mixed with tap water to suspend the nematodes. The soil slurry was passed through progressively finer mesh sieves (with sieve sizes mesh size 600, 250, 150, 75, 44 µm), 3 times for each sieve. The nematodes and some soil debris were caught on the sieves, and each time after sieving gently back-washed into a metal basin using a mister. Lastly, the content of the metal basin, which contained the nematodes, was poured over a cotton filter. Nematodes were identified to genus and genera were assigned to trophic groups (algal feeders, bacterivores, fungivores, plant-parasites, omnivores, and predators) according to Yeates et al. [29].

2.2.3 Vegetation. GPS coordinates were used to enable return to the same locations for sampling of herbaceous vegetation between 16 June and 25 August 2009. A flexible hoop encircling one square meter was placed on top of vegetation next to the GPS position where soil was gathered. The number of unique plant species was recorded, as well as the percentage of ground area covered by each of those species. Non-vegetative coverage was recorded as *leaf duff* or *bare soil*. Reference objects were held at arm's length to gauge percent coverage. Shrub data were only collected at the backslope, 10 m, and 50 m microtopographic areas because shrubs were not present at the shoulder, sideslope, or ditch, given the annual road maintenance. Shrubs that were within a 10 m² sweep were quantified and identified. A professional forester analyzed tree species along each transect at the 10 m and 50 m sites (Appendix 1). A 10x prism plot was used at each location to determine the basal area of the tree by species. The sites were then categorized using a Society of American Foresters classification system [19].

2.2.4 Road Attributes. Road attribute data (e.g., daily traffic, age of the road) were collected through database mining and interviews with Public Works officials [J. Cota personal communication, ^{31,31}]. Quantities of winter maintenance products applied to the road were determined by auditing the mean mass of product purchased for the past ten years to account for yearly weather differences and carryover of product from year to year. The average mass of product purchased was divided by the length of applicable roads in a maintenance jurisdiction to get a kilogram per meter estimation of chemical application. It is assumed for calculation purposes that all roads receive pickled sand, only two-lane paved roads and highways receive sodium chloride (NaCl), and only gravel roads receive calcium chloride (CaCl₂) dust suppressant. These products are not necessarily applied evenly across all miles of roads in the district; in practice they are applied as the conditions of the road warrant [J. Cota, personal communication]. However, for analysis purposes, we assume even product distribution.

2.3 Analytical Methodology

Nematode community composition was quantified using three complementary indices: 1) maturity which is a measure of ecological succession, 2) trophic diversity that measures the complexity of food webs, and 3) ratio of fungivorous to bacterivorous nematodes that reflects the dominant decomposition pathway. This suite of indices was chosen because they were complementary in characterizing the condition of soil food webs [18].

A 3-way analysis of variance (ANOVA) using the GLM procedure, with LSMEANS, utilizing type III sum of squares was used to determine whether soil chemistry, physical properties, or nematode communities varied by

topography, road type and road distance. Means comparisons of nematode samples were computed as orthogonal contrasts, and chemistry samples by a Tukey-Kramer Honestly Significant Difference (HSD). Data were analyzed using the statistical program SAS Version 9.3 (SAS Institute Inc., Cary, NC).

3. Results

The main ecological effects from roads on vegetation appears to be related to construction modifications required for a roadway (i.e., vegetation clearing and topography modification). The spatial extent of these modifications was correlated positively with road use intensity. Highways have the greatest ecological impact and gravel roads the least impact. The cleared area defined the type of plant community and the distance that road pollutants travel. Secondarily, road presence affected soil chemistry. Heavy metals (e.g., Pb, Cd, Cu, and Zn) correlated positively with road use intensity. In contrast, gravel roads have higher calcium content in nearby soil when compared to other road types (Fig. 3-1). Proximity to all road types made the soils more alkaline relative to the acidic soil of the adjacent native forest (Table 3-1).

Microtopography next to the road had marked effects on the composition of plant communities based on the direction of water flow. Ditch areas supported wetland plants, and had greater soil moisture and sulfur content, while plant communities closer to the road were characteristic of drier upland zones. The area beyond the edge of the forest did not appear to be affected chemically or physically by any of the road types, possibly due to the dense vegetation that typically develops outside of the managed right-of-way.

Soil chemistry and physical properties displayed a few general patterns (Table 3-2). One common pattern was higher values closer to the road and with increasing road use intensity class. This included lead (Pb), which had greatest mean values near the shoulder and in the case of highway roads (Fig. 3-1). Spatial patterns of zinc (Zn) and copper (Cu) were similar to Pb, with concentrations greatest near the roadside and with most intense road use. A second common pattern was higher values at greater distances from the road which included organic matter, cation exchange capacity and nutrients.




Figure 3-1. Mean levels of lead (mg/kg) in roadside soil based upon categorical distance from road. Error bars represent standard error. Paved refers to 2-lane paved roads. 10M is 10 meters into the forest from the forest edge, and 50M is 50 meters from the road crown.

A second group of measurements also had higher values near the road, but with highest values in the case of gravel roads, and lowest in the case of highways (Table 3-1). Mean calcium (Ca) values were greatest in gravel road samples near the roadside, with 2-lane paved roads being second highest, and highway samples smallest (Fig. 3-3). Calcium varied by road type, distance and topography, as well as interaction term of road and distance (all cases $p < 0.0001$).

Similar to Ca, pH exhibited higher means near the shoulder with gravel roads, however the differences between the road types was not as pronounced. As expected, gravel roads also had the highest amount of rocky debris (particles $> 2\text{mm}$), particularly in cases of the sideslope and ditch. Mean bulk density was greatest near the roadside and decreased with distance. There was an interaction by road type and topography with 2-lane paved and highway alternating as the smallest means, but the overall trend showed higher gravel bulk density in the sideslope, ditch and backslope, and convergence at 50m. Percent clay was also highest in the case of gravel roads, particularly at the shoulder and in case of up-sloping transects.





Figure 3-2. Mean levels of calcium (mg/kg) in roadside soil based upon categorical distance from road. Error bars represent standard error. Paved refers to 2-lane paved roads. 10M is 10 meters into the forest from the forest edge, and 50M is 50 meters from the road.

Table 3-1. Soil properties for each of three road types that correspond with increase travel intensity: gravel, two-laned paved, and highway.

Soil Component	Road Type 1	>=<	Road Type 2	p-value
% Organic Matter	Gravel	<	Two-Lane Paved	0.0304
	Two-Lane Paved	=	Highway	0.2553
	Highway	>	Gravel	0.0001
Nickel	Gravel	=	Two-Lane Paved	0.2898
	Two-Lane Paved	<	Highway	0.0001
	Highway	>	Gravel	0.0001
% Potassium	Gravel	=	Two-Lane Paved	0.0752
	Two-Lane Paved	<	Highway	0.0484
	Highway	>	Gravel	0.0001
% Magnesium	Gravel	=	Two-Lane Paved	0.1005
	Two-Lane Paved	<	Highway	0.0001
	Highway	>	Gravel	0.0001
Calcium	Gravel	>	Two-Lane Paved	0.0001
	Two-Lane Paved	=	Highway	0.0769
	Highway	<	Gravel	0.0001
Manganese	Gravel	>	Two-Lane Paved	0.0001
	Two-Lane Paved	=	Highway	0.9981
	Highway	<	Gravel	0.0001

Table 3-2. Mean values of soil component from all road types separated into components that decrease with distance from the road, and those that increase with distance from the road. Metrics not listed had a less defined gradient



Sodium concentration, as well as electrical conductivity (EC), exhibited a unique pattern. Sodium (Na) was highest in 2-lane paved soil samples, specifically at the ditch and backslope (Fig. 3-3).




Figure 3-3. Mean levels of sodium (mg/kg) in roadside soil based upon categorical distance from road. Error bars represent standard error. Paved refers to 2-lane paved roads. 10M is 10 meters into the forest from the forest edge, and 50M is 50 meters from the road.

Nematodes. The nematode community proved to be a useful bioindicator of soil condition. A total of 119 genera were identified in the entire data set (Appendix 2). The range of samples contained between 0 and 37 genera, while the average number of genera across all samples (richness) was 18.8. Over the entire data set there was only one algal feeder found, 36 bacterivore genera, 21 fungivore genera, 29 plant-parasitic genera, 11 omnivore genera, 20 predator genera. Values of the MI were greater for highway than paved or gravel regardless of distance from the road (Fig 3-4). Values were lower near than further from the road




Figure 3-4. Mean Maturity index of nematode communities in roadside soil based upon categorical distance from road. Error bars represent standard error. Paved refers to 2-lane paved roads. 10M is 10 meters into the forest from the forest edge, and 50M is 50 meters from the road.

Reflective of the trends with bacterivore and fungivore fractions, the overall trend in the $F:F+B$ ratio was least nearest the roadside and increased with distance from the road (Figure 3-5). Interestingly, the 2-lane paved roads had consistently lower ratios than both gravel and highway roads. The means were significantly different by road type and distance ($p < 0.0001$).




Figure 3-5. Mean ratio of fungal feeding to fungal and bacterial feeding nematodes in roadside soil based upon categorical distance from road. Error bars represent standard error. Paved refers to 2-lane paved roads. 10M is 10 meters into the forest from the forest edge, and 50M is 50 meters from the road.\

4. Implementation/Tech Transfer

This research is part of the University of Vermont Transportation Center Signature Project entitled “Integrated Land-use, Transportation and Environmental Modeling: Complex Systems Approaches and Advanced Policy Applications”, into which the results of this research will eventually be incorporated (Fig. 4-1). The signature project has brought together research from both transportation and environmental science, with the intention of integrating research results into new models. These models will be created by enhancing existing traffic modeling software (UrbanSim and TRANSIMS). This project was developed with the aim of being a national showcase for the testing and validation of integrated models and for more sustainable transportation planning.

Transportation Research Center Signature Project 1




Figure 4-1. Schema of projects integrated together under TRC Signature 1. This study, 1F, is highlighted.

The results of the primary objective were presented at the UVM Student Research Conference:

- 2009- Spatial extent of the impact of transported road materials on the ecological function of forested landscapes, Kristin Williams
- 2012 - Spatial effects of roads on soil nematode communities in forested areas of Vermont, Kristin Williams and David Asmussen

Two M.S. theses were based on analysis of the data gathered in this project:

- The spatial effects of road use intensity on forest plant communities and soil chemistry: does the road less traveled by, make all the difference? – Dave Asmussen, 2010
- Spatial effects of roads on soil nematode communities in forested areas of Vermont – Kristin Williams, 2012

5. Conclusions

The results of both abiotic and biotic soil results present a complex picture of roadside soil ecology in Vermont. These results did suggest the accumulation of pollutants near the roadside, and the combined effects of these pollutants and habitat alterations of the roadside are associated with changes in the soil nematode community, based upon both distance from the roadside and road type. Results suggest that pollutants and most of the ecological effects are concentrated near the roadside, and with higher intensity roads. However,

each road type had a unique effect with different pollutants being more prominent.

The roadside grass-verge community was substantially different from the forest in terms of soil nematodes, and plant habitat is probably an influencing factor in these results in addition to chemical and physical soil alteration. Ultimately roadside effects are linked to road type due to the nature of road building, such that highway roads have a much broader roadside verge, 2-lane paved roads a moderate size road verge, and gravel roads a relatively narrow roadside verge. Specific local roadside conditions influence this distance and considerations of local conditions may help designers reduce road effects on the surrounding landscape.

This research supports findings that within an established forest environment, the use-intensity effects of roads on soil and vegetation are contained within the corresponding right-of-way. Increased traffic contributes to higher levels of heavy metals in roadside soil, but these effects do not extend past the forest edge. However, more traffic is correlated positively with wider maintained roadside areas, so the forest edge is effectively farther back compared to lesser-used roads.

Maintained roadside area width is the most significant factor responsible for the roadside environment's deviation from native plant and soil composition. Large maintained areas close to the road are more similar to grassland communities than they are to native forest communities. Plants at the edge of the forest grow in response to increased resource availability and buffer the interior forest from roadside effects.

To reduce impacts of roads, the width of cleared area should be reduced and traffic consolidated to fewer individual vehicles on the road. The forest edge should be cultivated to maximize the insulating effect and maintain low-resource-adapted native plant communities within the forest. By reducing the transportation system's physical footprint and cultivating native vegetation borders, we can maintain natural plant communities and stem the introduction of chemicals into the environment.

6. References (in order of citation)

1. Trombulak, S. C. and Frissell, C. A. 2000. "Review of Ecological Effects of Roads on Terrestrial and Aquatic Communities". *Conservation Biology* Vol. 14, pp. 18-30.
2. Spooner, P. G. and Smallbone, L. 2008. "Effects of Road Age on the Structure of Roadside Vegetation in South-Eastern Australia". *Agriculture, Ecosystems and Environment* Vol. 129, pp. 57-64.
3. Cilliers, S. S. and Bredenkamp, G. J. 1999. "Vegetation of Road Verges on an Urbanization Gradient". *Landscape and Urban Planning* Vol. 46, pp. 217-239.
4. Wardle, D. A., Bardgett, R. D., Klironomos, J. N., Setälä, H., van der Putten, W. H. and Wall, D. H. 2004. "Ecological Linkages between Aboveground and Belowground Biota". *Science* Vol. 304, pp. 1629-1633.
5. Sullivan, J. J., Williams, P. A., Timmins, S. M. and Smale, M. C. 2009. "Distribution and Spread of Environmental Weeds along New Zealand Roadsides". *New Zealand Journal of Ecology* Vol. 33, pp. 190-204.
6. Forman, R. T. T. and Deblinger, R. D. 2000. "The Ecological Road-Effect Zone of a Massachusetts (USA) Suburban Highway". *Conservation Biology* Vol. 14, pp. 36-46.
7. Gobel, P., Dierkes, C. and Coldewey, W. G. 2006. "Storm Water Runoff Concentration Matrix for Urban Areas". *Journal of Contaminant Hydrology* Vol. 91, pp. 26-42.
8. Jodoin, Y., Lavoie, C., Villeneuve, P., Theriault, M., Beaulieu, J. and Belzile, F. 2008. Highways as Corridors and Habitats for the Invasive Common Reed *Phragmites australis* in Quebec, Canada. *Journal of Applied Ecology* Vol. 45, pp. 459-466.
9. Auerbach, N. A., Walker, M. D. and Walkerm D. A. 1997. "Effects of Roadside Disturbance on Substrate and Vegetation Properties in Arctic Tundra". *Ecological Applications* Vol. 7, pp. 218-235.
10. Grime, J. P. 1977. "Evidence for the Existence of Three Primary Strategies in Plants and its Relevance to Ecological and Evolutionary Theory". *The American Naturalist* Vol. 111, pp. 1169-1194.
11. Sahlodin, A. M., Sotudeh-Gharebagh, R. and Zhu Y. 2007. Modeling of Dispersion Near Roadways Based on the Vehicle-Induced Turbulence Concept. *Atmospheric Environment* Vol. 41, pp. 92-102.
12. Venner, M. 2006. Control of Invasive Species: A Synthesis of Highway Practice. Transportation Research Board, Washington D.C., 115p.
13. Cape, J. N., Tang, Y. S., van Dijk, N., Love, L., Sutton, M. A. and Palmer, S. C. F. 2004. "Concentrations of Ammonia and Nitrogen Dioxide at Roadside Verges, and their Contribution to Nitrogen Deposition". *Environmental Pollution* Vol. 132, pp. 469-478.
14. Azizian, M. F., Nelson, P. O., Thayumanavan, P. and Williamson, K. J. 2003. "Environmental Impact of Highway Construction and Repair Materials on Surface And Ground Waters: Case Study: Crumb Rubber Asphalt

- Concrete. *Waste Management* Vol. 23, pp. 719-728.
15. Forman, R. T. T., Sperling, D., Bissonette, J. A., Clevenger, A. P., Cutshall, C. D., Dale, V. H., Fahrig, L., France, R., Goldman, C. R., Heanue, K., Jones, J. A., Swanson, F. J., Turrentine, T. and Winter, T. C. 2003. *Road Ecology, Science and Solutions*. Island Press, Washington, DC, 481p.
 16. Adachi, K. and Tainosho, Y. 2004. "Characterization of Heavy Metal Particles Embedded in Tire Dust". *Environment International* Vol. 30, pp. 1009-1017.
 17. United States Department of Transportation, Federal Highway Administration. 2011. Flexibility in Highway Design. Available at: <http://www.fhwa.dot.gov/environment/flex/ch03.htm> (accessed October 27, 2011). U.S. Department of Transportation, Washington, DC
 18. Neher, D. A. 2001. "Role of Nematodes in Soil Health and Their Use As Indicators". *Journal of Nematology* Vol. 33, pp. 161-168.
 19. Eyre, F. H. 1980. *Forest Cover Types of the United States and Canada*. The Society of American Foresters, 148p.
 20. Thompson, E. H. and Sorenson, E. R. 2005. *Wetland, Woodland, Wildland*. The Nature Conservancy, Lebanon, NH, 456p.
 21. Fitzgerald, E. P. and Parker, S. P. 2009. Muddy Brook phase 1 and 2 stream geomorphic assessment summary. p. 1-26. In Fitzgerald Environmental Associates, LLC. Vermont River Management Program.
 22. Haahr, M. 2009. Random.Org. Available at www.random.org (accessed 2/10/2009). TSDA, Dublin.
 23. Karim, M. N. and Mallik A.U. 2008. "Roadside Revegetation by Native Plants: I. Roadside Microhabitats, Floristic Zonation and Species Traits". *Ecological Engineering* Vol. 32, pp. 222-237.
 24. Morgan, M. F. 1941. *Chemical Soil Diagnosis by the Universal Soil Testing System*. Connecticut Agricultural Experiment Station Bulletin. Connecticut.
 25. Lunt, H. A., Swanson, C. L. and Jacobson, H. G. M. 1950. *The Morgan Soil Testing System*. Connecticut Agricultural Experiment Station Bulletin. Connecticut.
 26. Walkley, A. and Black, I. A. 1934. "An Examination of Degtjareff Method for Determining Soil Organic Matter and a Proposed Modification of the Chromic Acid Titration Method". *Soil Science* Vol. 37, pp. 29-37.
 27. Whitehead, A. G. and Hemming, J. R. 1965. "A Comparison of Some Quantitative Methods of Extracting Small Vermiform Nematodes from Soil". *Annals of Applied Biology* vol. 55, pp. 25-38.
 28. s' Jacob, J. J. and van Bezooijen, J. 1984. *A Manual for Practical Work in Nematology*. Wageningen Agricultural University, Wageningen, The Netherlands.
 29. Yeates, G. W., Bongers, T., de Goede, R. G. M., Freckman, D. W. and Georgieva, S. S. 1993. "Feeding Habits in Soil Nematode Families and Genera – An Outline for Soil Ecologists". *Journal of Nematology* vol. 25, pp. 315-331.
 30. United States Department of Transportation, 2010. *Highway Statistical*

- Publications, Federal Highway Administration. Available at:
<http://www.fhwa.dot.gov/policy/ohpi/hss/hsspubs.cfm> (accessed June 30, 2010). U.S. Department of Transportation, Washington, DC.
31. Vermont Agency of Transportation, Traffic Research Unit, June 2009. Automatic Traffic Recorder Station History 1999-2008, Available at:
<http://www.aot.state.vt.us/planning/documents/trafresearch/publications/pub.htm>, Accessed on: 3/23/2010.
32. Magee, D. W. and Ahles, H. E. Flora of the Northeast, A Manual of the Vascular Flora of New England and Adjacent New York. 2nd Edition. University of Massachusetts Press, Amherst, 1214p.

7. Appendix 1.

List of plant species and their coverage for road type and distance.
 Nomenclature follows Magee and Ahles, [32].

Family	Genus	Species	Gravel Shoulder	Gravel Sideslope	Gravel Ditch	Gravel Backslope	Gravel 10m	Gravel 50m	Paved Shoulder	Paved Sideslope	Paved Ditch	Paved Backslope	Paved 10m	Paved 50m	Highway Shoulder	Highway Sideslope	Highway Ditch	Highway Backslope	Highway 10m	Highway 50m
Aceraceae	Acer	saccharum																		
Acoraceae	Acorus	calamus																		
Alismataceae	Sagittaria	latifolia																		
Alliaceae	Allium	triccoccum																		
Anacardiaceae	Rhus	typhina																		
	Toxicodendron	rydbergii																		
Apiaceae	Daucus	carota																		
	Pastinaca	sativa																		
	Sium	suave																		
	Zizia	aurea																		
Aquifoliaceae	Ilex	verticillata																		
Araceae	Arisaema	triphillum																		
Araliaceae	Aralia	nudicaulis																		
Asclepiadaceae	Asclepias	syriaca																		
Asteraceae	Achillea	millefolium																		
	Ambrosia	artemisiifolia																		
	Arcticum	minus																		
	Artemisia	vulgaris																		
	Aster	cordifolius																		
	Aster	lateriflorus																		
	Aster	sp.																		
	Bidens	frondosa																		
	Cichorium	intybus																		
	Cirsium	vulgare																		
	Conyza	canadensis																		
	Erigeron	philadelphicus																		
	Eupatorium	maculatum																		
	Euthamia	graminifolia																		
	Hieracium	caespitosum																		
	Hieracium	sp.																		
	Leucanthemum	vulgare																		
	Matricaria	discoidea																		
	Prenanthes	alba																		
	Prenanthes	trifoliolata																		
	Rudbeckia	hirta																		
	Solidago	caesia																		
	Solidago	canadensis																		
	Solidago	flexicaulis																		
	Solidago	gigantea																		
	Solidago	rugosa																		
	Sonchus	oleraceus																		
	Taraxacum	officinale																		
	Tussilago	farfara																		
Balsaminaceae	Impatiens	capensis																		
Berberidaceae	Caulophyllum	thalictroides																		
Betulaceae	Ostrya	virginiana																		
Caprifoliaceae	Lonicera	morrowii																		
	Sambucus	canadensis																		
	Viburnum	acerifolium																		
	Viburnum	dentatum																		
	Viburnum	lentago																		
Caryophyllaceae	Silene	vulgaris																		
	Stellaria	graminea																		
Chenopodiaceae	Chenopodium	album																		
	Chenopodium	sp.																		

Family	Genus	Species	Gravel Shoulder	Gravel Sideslope	Gravel Ditch	Gravel Backslope	Gravel 10m	Gravel 50m	Paved Shoulder	Paved Sideslope	Paved Ditch	Paved Backslope	Paved 10m	Paved 50m	Highway Shoulder	Highway Sideslope	Highway Ditch	Highway Backslope	Highway 10m	Highway 50m
Clusiaceae	Hypericum	perforatum																		
Convallariaceae	Clintonia	borealis																		
	Maianthemum	canadense																		
	Polygonatum	pubescens																		
	Smilacina	racemosa																		
Convolvulaceae	Calystegia	sepium																		
Cornaceae	Cornus	amomum																		
	Cornus	racemosa																		
	Cornus	sericea																		
Cupressaceae	Thuja	occidentalis																		
Cyperaceae	Carex	arctata																		
	Carex	gracillima																		
	Carex	grayi																		
	Carex	gynandra																		
	Carex	hystericina																		
	Carex	intumescens																		
	Carex	ovalis																		
	Carex	plantaginea																		
	Carex	rosea																		
	Carex	sp.																		
	Carex	swanii																		
	Carex	vulpinoidea																		
	Eleocharis	tenuis																		
	Scirpus	atrovirens																		
	Scirpus	microcarpus																		
	UID	UID																		
	Dennstaedtiaceae	Pteridium	aquilinum																	
Dryopteridaceae	Athyrium	filix-femina																		
	Cystopteris	bulbifera																		
	Deparia	acrostichoides																		
	Dryopteris	intermedia																		
	Dryopteris	marginalis																		
	Matteuccia	struthiopteris																		
	Onoclea	sensibilis																		
	Polystichum	acrostichoides																		
Equisetaceae	Equisetum	arvense																		
	Equisetum	hyemale																		
Ericaceae	Gaultheria	procumbens																		
	Vaccinium	corymbosum																		
Fabaceae	Amphicarpaea	bracteata																		
	Lotus	corniculatus																		
	Medicago	lupulina																		
	Melilotus	alba																		
	Trifolium	arvense																		
	Trifolium	pratense																		
	Trifolium	repens																		
Vicia	cracca																			
Gentianaceae	Centaurium	erythraea																		
Geraniaceae	Geranium	robertianum																		
Grossulariaceae	Ribes	cynosbati																		
Juncaceae	Juncus	effusus																		
	Juncus	sp.																		
Lamiaceae	Glechoma	hederacea																		
	Prunella	vulgaris																		
Lycopodiaceae	Lycopodium	obscurum																		

Family	Genus	Species	Gravel Shoulder	Gravel Sideslope	Gravel Ditch	Gravel Backslope	Gravel 10m	Gravel 50m	Paved Shoulder	Paved Sideslope	Paved Ditch	Paved Backslope	Paved 10m	Paved 50m	Highway Shoulder	Highway Sideslope	Highway Ditch	Highway Backslope	Highway 10m	Highway 50m
Lythraceae	Lythrum	salicaria																		
Melanthiaceae	Veratrum	viride																		
Monotropaceae	Monotropa	uniflora																		
Nyctaginaceae	Mirabilis	nyctaginea																		
Oleaceae	Fraxinus	americana																		
Onagraceae	Circaea	lutetiana																		
	Oenothera	biennis																		
Orchidaceae	Epipactis	helleborine																		
Osmundaceae	Osmunda	cinnamomea																		
	Osmunda	claytoniana																		
	Osmunda	regalis																		
Oxalidaceae	Oxalis	europaea																		
	Oxalis	stricta																		
Plantaginaceae	Plantago	major																		
Poaceae	Bromus	inermis																		
	Carex	hirtifolia																		
	Dactylis	glomerata																		
	Echinochloa	sp.																		
	Elymus	repens																		
	Festuca	sp.																		
	Glyceria	striata																		
	Phalaris	arundinacea																		
	Phleum	pratense																		
	Setaria	glauca																		
	Setaria	pumila																		
	UID	UID																		
Polygonaceae	Polygonum	aviculare																		
	Polygonum	persicaria																		
	Polygonum	sagittatum																		
	Rumex	crispus																		
Polypodiaceae	Polypodium	virginianum																		
Primulaceae	Lysimachia	ciliata																		
	Lysimachia	nummularia																		
	Lysimachia	terrestris																		
	Trientalis	borealis																		
Pyrolaceae	Moneses	uniflora																		
Ranunculaceae	Anemone	virginiana																		
	Aquilegia	canadensis																		
	Caltha	palustris																		
	Ranunculus	acris																		
	Thalictrum	dioicum																		
Rhamnaceae	Rhamnus	cathartica																		
Rosaceae	Fragaria	virginiana																		
	Geum	canadense																		
	Prunus	virginiana																		
	Rosa	multiflora																		
	Rubus	alleggheniensis																		
	Rubus	idaeus																		
	Rubus	odoratus																		
	Rubus	pubescens																		
	Spiraea	alba																		
Rubiaceae	Galium	palustre																		
	Galium	triflorum																		
	Galium	mollugo																		
	Mitchella	repens																		

Family	Genus	Species	Gravel Shoulder	Gravel Sideslope	Gravel Ditch	Gravel Backslope	Gravel 10m	Gravel 50m	Paved Shoulder	Paved Sideslope	Paved Ditch	Paved Backslope	Paved 10m	Paved 50m	Highway Shoulder	Highway Sideslope	Highway Ditch	Highway Backslope	Highway 10m	Highway 50m
Salicaceae	Populus	tremuloides																		
	Salix	nigra																		
Saxifragaceae	Tiarella	cordifolia																		
Scrophulariaceae	Chelone	glabra																		
	Verbascum	thapsus																		
	Veronica	officinalis																		
Solanaceae	Solanum	dulcamara																		
Thelypteridaceae	Thelypteris	noveboracensis																		
	Thelypteris	palustris																		
Trilliaceae	Trillium	erectum																		
	Trillium	undulatum																		
Typhaceae	Typha	latifolia																		
Urticaceae	Pilea	pumila																		
Verbenaceae	Verbena	hastata																		
Violaceae	Viola	sp.																		
Vitaceae	Parthenocissus	quinquefolia																		
	Vitis	riparia																		
Unknown Forb #1	UID	UID																		
Unknown Graminoid #1	UID	UID																		
Unknown Graminoid #2	UID	UID																		
Unknown Graminoid #3	UID	UID																		
Other Ground Cover																				
Bare Soil																				
Leaf Duff																				

Appendix 2.

Classification of nematode genera by c-p value of the Maturity Index and trophic group, and the percentage of samples with nonzero abundance of specified genera by road type.

Genus	Family	c-p	Trophic	Road Type		
				Gravel ^a	2-laned paved ^b	highway ^a
<i>Achromadora</i> ^d	Achromadoridae	3	A	42	50	50
<i>Acrobeles</i>	Cephalobidae	2	B	7	19	12
<i>Acrobeloides</i>	Cephalobidae	2	B	77	83	72
<i>Ailaimus</i> ^{c,d}	Alaimidae	4	B	58	31	38
<i>Anaplectus</i> ^d	Plectidae	2	B	20	17	10
<i>Anatonchus</i>	Anatonchidae	4	P	0	0	3
<i>Anomyctus</i>	Aphelenchoididae	2	B	18	19	3
<i>Aphelechoides</i>	Aphelenchoididae	2	F	87	84	85
<i>Aphelenchus</i> ^{c,d}	Aphelenchidae	2	F	40	40	57
<i>Aporcelaimellus</i>	Aporcelaimidae	5	P	15	7	12
<i>Aporcelaimium</i>	Aporcelaimidae	5	O	0	0	2
<i>Aporcelaimus</i>	Aporcelaimidae	5	P	8	7	7
<i>Axonchium</i>	Belonidiridae	5	PP	3	3	8
<i>Basiria</i>	Tylenchidea	2	F	3	5	8
<i>Bastiania</i>	Bastianiidae	3	B	8	9	0
<i>Bitylenchus</i>	Dolichoridae	3	PP	0	5	8
<i>Boleodorus</i> ^c	Tylenchidae	2	F	57	31	50
<i>Bunonema</i>	Bunonnematidae	1	B	3	0	0
<i>Bursaphelenchus</i>	Aphelenchoididae	2	F	0	0	2
<i>Carcharolaimus</i>	Discolaimidae	5	P	0	0	2
<i>Cephalenchus</i> ^d	Tylodoridae	2	PP	12	10	12
<i>Cephalobus</i>	Cephalobidae	2	B	87	88	68
<i>Cervidellus</i> ^c	Cephalobidae	2	B	38	31	17
<i>Chiloplacus</i>	Cephalobidae	2	B	7	3	8
<i>Chronogaster</i>	Leptolaimidae	2	B	2	7	0
<i>Clarkus</i> ^d	Monochidae	4	P	30	19	17
<i>Coomansis</i>	Monochidae	4	P	0	3	2
<i>Coslenchus</i>	Tylenchidae	2	F	6.7	16	17
<i>Criconema</i>	Criconematidae	3	PP	0	0	2
<i>Criconemella</i>	Criconematidae	3	PP	2	0	5
<i>Criconemoides</i>	Criconematidae	3	PP	2	0	2
<i>Croossonema</i>	Criconematidae	3	PP	0	7	5
<i>Cylindrolaimus</i>	Diphtherophoridae	3	B	0	2	0
<i>Dactyluraxonchium</i>	Belonidiridae	5	PP	2	0	2
<i>Diphtherophora</i>	Diphtherophoridae	3	F	17	17	18

<i>Diplogaster</i>	Diplogasteroididea	1	B	7	0	0
<i>Diplogasteritus</i>	Diplogasteroididea	1	B	0	2	0
<i>Discolaimus</i>	Discolaimidae	5	P	2	0	2
<i>Ditylenchus</i>	Anguinidae	2	F	62	55	65
<i>Dorydorella</i>	Qudsianematidae	4	O	3	0	0
<i>Dorylaimellus</i>	Belonidiridae	5	F	5	2	3
<i>Dorylaimoides</i>	Leptonchidae	4	O	0	0	2
<i>Dorylaimus</i>	Dorylaimidae	4	O	2	0	0
<i>Echphyadophoroides</i>	Tylenchidae	2	F	0	0	2
<i>Echphyadophorus</i>	Tylenchidae	2	F	0	2	0
<i>Enchodellus</i>	Nordiidae	4	O	0	2	0
<i>Epidorylaimus</i>	Qudsianematidae	4	O	23	5	7
<i>Etamphidelus</i>	Alaimidae	4	B	0	3	0
<i>Eucephalobus</i>	Cephalobidae	2	B	92	86	83
<i>Eudorylaimus</i>	Qudsianematidae	4	O	32	29	18
<i>Eumonhystera</i>	Monhysteridae	2	B	80	84	80
<i>Euteratocephalus</i>	Teratocephalidae	3	B	0	0	3
<i>Filenchus</i>	Tylenchidae	2	F	93	88	93
<i>Fraglenchus</i>	Tylenchidae	2	F	0	0	2
<i>Geomonhystera</i>	Monhysteridae	2	B	0	0	2
<i>Helicotylenchus</i> ^c	Hoplolaimidae	3	PP	43	41	70
<i>Hemicycliphora</i> ^d	Hemicyliophoridae	3	PP	13	19	10
<i>Heterocephalobus</i> ^d	Cephalobidae	2	B	15	7	10
<i>Heterodera</i>	Heteroderidae	3	PP	0	2	0
<i>Ironus</i>	Ironidae	4	P	0	0	2
<i>Lelenchus</i>	Tylenchidae	2	F	0	10	7
<i>Leptolaimus</i>	Leptolaimidae	2	B	17	5	3
<i>Leptonchus</i>	Leptonchidae	4	F	3	0	2
<i>Longidorus</i>	Longidoridae	5	PP	8	2	2
<i>Loofia</i>	Hemicyliophoridae	3	PP	2	2	0
<i>Macroposthnia</i> ^d	Criconematidae	3	PP	8	9	17
<i>Malenchus</i>	Tylenchidae	2	F	27	21	18
<i>Meloidogyne</i>	Meloidogynidae	3	PP	3	9	10
<i>Merlinus</i>	Dolichoridae	3	PP	0	2	0
<i>Mesodorylaimus</i> ^d	Thornenematidae	5	O	18	5	13
<i>Metateratocephalus</i>	Teratocephalidae	3	B	3	7	0
<i>Miconchus</i>	Anatonchidae	4	P	0	0	2
<i>Microdorylaimus</i>	Qudsianematidae	4	O	5	0	0
<i>Monhystera</i> ^d	Monhysteridae	2	B	15	17	17
<i>Mononchoides</i>	Neodiplogasteroidid	1	B	2	0	0
<i>Mononchus</i>	Monochidae	4	P	7	0	5
<i>Mylonchulus</i>	Monochidae	4	P	20	31	13
<i>Neopsilenchus</i>	Tylenchidae	2	F	13	14	60
<i>Nygolaimus</i>	Nygolaimidae	5	P	2	0	2
<i>Ogma</i>	Criconematidae	3	PP	0	2	2

<i>Oxydirus</i>	Belondiridae	5	PP	2	2	5
<i>Panagrobellus</i>	Panagrolaimidae	1	B	0	5	2
<i>Panagrolaimus</i> ^d	Panagrolaimidae	1	B	20	24	22
<i>Paractinolaimus</i>	Actinolaimidae	5	P	0	0	2
<i>Paramphidellus</i> ^{c,d}	Alaimidae	4	B	42	32	52
<i>Paraplectonema</i>	Leptonchidae	4	B	0	0	2
<i>Paratrichodoros</i>	Trichodoridae	4	PP	0	2	0
<i>Paratylenchus</i> ^d	Paratylenchidae	2	PP	15	24	17
<i>Paravulvulus</i>	Nygolaimidae	5	P	2	0	0
<i>Paraxonchium</i>	Aporcelaimidae	5	P	2	0	0
<i>Plectus</i>	Plectidae	2	B	92	84	85
<i>Pleurotylenchus</i>	Tylodoridae	2	PP	0	5	7
<i>Pratylenchoides</i>	Pratylenchidae	3	PP	0	2	2
<i>Pratylenchus</i> ^c	Pratylenchidae	3	PP	15	29	35
<i>Prismatolaimus</i>	Prismatolaimidae	3	B	35	50	37
<i>Pristionchus</i>	Neodiplogasteroidid	1	B	0	3	2
<i>Prodorylaimus</i> ^c	Thornenematidae	5	O	45	24	47
<i>Protodiplogasteroides</i>	Diplogasteroididea	1	B	2	0	0
<i>Psilenchus</i>	Psilenchidea	2	PP	10	3	15
<i>Pungentus</i>	Nordiidae	4	PP	17	28	18
Rhabditidae ^{+d}	Rhabditidae	1	PP	30	38	37
<i>Rhabdolaimus</i> ^{c,d}	Rhabdolaimidae	3	B	15	21	38
<i>Rotylenchus</i> ^d	Hoplolaimidae	3	PP	13	16	5
<i>Scutylenchus</i>	Dolichoridae	3	PP	0	0	5
<i>Seinura</i>	Aphelenchoididae	2	P	0	5	5
<i>Stenonchulus</i>	Onchulidae	3	P	2	0	0
<i>Teratocephalus</i>	Teratocephalidae	3	B	43	52	32
<i>Thonus</i> ^d	Qudsianematidae	4	O	48	41	30
<i>Tobrilus</i>	Tobrilidae	3	P	2	2	0
<i>Triplya</i>	Triplylidae	3	P	0	0	2
<i>Trischistoma</i>	Triplylidae	3	P	0	0	2
<i>Tylencholaimellus</i>	Leptonchidae	4	F	8	9	22
<i>Tylencholaimus</i>	Leptonchidae	4	F	50	45	52
<i>Tylenchus</i>	Tylenchidae	2	F	83	72	73
<i>Tyocephalus</i>	Plectidae	2	B	7	12	2
<i>Wilsonema</i>	Plectidae	2	B	17	33	17
<i>Xiphinema</i> ^d	Longidoridae	5	PP	13	9	5
<i>Zanenchus</i>	Tylenchidae	2	F	0	0	2

a. $n = 60$, b. $n = 58$,

chi squared test $p < 0.05$: c. for road type; d. for topography, percent nonzero not listed.

Trophic Groups defined as: A. Algal Feeder, B. Bacterial Feeder, F. Fungal Feeder,

O. Omnivores, P. Predators, PP. Plant (Root) Feeders

road type $df=2$, topography $df=1$, + Rhabditidae only identified to family.

8. Equations

Equation 2. Nematode maturity index

$MI = \sum [(v_i f_i) / n]$ where v_i equals the *c-p* value of the *i*th family, f_i equals the frequency of the *i*th family in the sample, and n equals the total number of individual nematodes in a sample.

Equation 3. Nematode trophic diversity index

Hills $N1 = \exp [- \sum P_i (\ln P_i)]$ where P_i is the proportion of trophic group *i*.

Equation 4. Ratio of fungivorous nematodes (FN) to bacterivorous nematodes (BN)

$F:F+B = (no. FN) / (no. FN + no. BN) \times 100$.