

# Closing the Nutrient Loop Through an Innovative Organic Fertilizer Technology Field Tested for Beet Vegetable Crop

A new byproduct of wastewater treatment can provide an organic alternative to chemical fertilizers while improving water quality.

## Nutrient Pollution and Harmful Algae Blooms

Excess nutrients from agricultural, industrial, and wastewater runoff result in harmful algae blooms in rivers, lakes, and the ocean. Inorganic nitrogen and phosphorous can fuel the rapid growth of algae, which deprives the aquatic environment of oxygen and sunlight. This process, eutrophication, results in anoxic “dead zones” and threatens ecosystems and access to clean water.


In agriculture, this nutrient pollution can be controlled by implementing and improving practices that reduce runoff, such as building good soil structure and not over-fertilizing nitrogen and phosphorus using slow release fertilizers.



Algal bloom in Lake Erie, 2017. Image from NOAA.

## Wastewater Treatment

Currently, phosphorus usage efficiency at farms is less than 20%, with the rest ending up in wastewater and surface waters. The waste to value technology developed by GSR Solutions can be used as a nutrient treatment system to remove excessive nutrients from both the digested or non-digested waste effluent streams and convert them into specialized microbial biomass, which prevents them from being flushed back out into rivers and lakes. Byproducts of this technology can be repurposed for fertilizers, food, or biofuel. Recycling nutrients through the process returns nutrients to crops, improves water quality, and reduces nutrient loss by “closing the loop”.



How the novel bio-based process for valued products works.

## From Nutrient Recovery to Organic Fertilizer

Using organic fertilizers helps to reduce nutrient runoff in comparison to synthetic fertilizers, resulting in improved water quality. Organic fertilizers increase organic matter, build good soil structure/aggregate stability, and increase the soil’s ability to retain water and nutrients. Organic fertilizers also release nutrients slowly as they decompose, making it harder to over-fertilize. This bio-based technology can provide an organic alternative to chemical fertilizers.

The advanced microbial-based biomass in the waste to value process is relatively quick to grow, as it captures nutrients from organic waste streams. Implementation of this process at the origin of nutrient runoff for fertilizer production can help to repair an excess algae problem down the watershed and improve water quality and quality of life, especially in rural areas.

## NORTHWEST CROPS & SOILS PROGRAM



Northwest Crops and Soils Program | 278 So. Main Street, Suite 2 | St. Albans, VT 05478

802-524-6501 or 1-800-639-2130 (toll-free in Vt.) | [susan.brouillette@uvm.edu](mailto:susan.brouillette@uvm.edu)

[uvm.edu/extension/nwcrops](http://uvm.edu/extension/nwcrops)

Sources: <https://oceanservice.noaa.gov/facts/hab-solutions.html>



## CULTIVATING HEALTHY COMMUNITIES

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. University of Vermont Extension, Burlington, Vermont. University of Vermont Extension, and U.S. Department of Agriculture, cooperating, offer education and employment to everyone without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or familial status. Any reference to commercial products, trade names, or brand names is for information only, and no endorsement or approval is intended.

COLLEGE OF AGRICULTURE AND LIFE SCIENCES

# Closing the Nutrient Loop Through an Innovative Organic Fertilizer Technology Field Tested for Table Beets


## Results From Field Trials

In May 2022, UVM Extension's Northwest Crop and Soils Team trialed a couple of nutrient-recovered fertilizer made from the treatment of anaerobically digested liquid dairy manure effluent with the Ag-biotechnology developed by GSR Solutions in comparison to commercial chemical and organic fertilizers. The GSR fertilizers (3% N) and (4% N) were compared to North Country Organic Pro-Gro (5% N), soybean meal (6% N), hemp seed meal (5% N), urea (46% N), and an untreated control.

**There were no statistical differences between Beets yield as well as soil quality between the four treatments.**

Compared to the Pro-Gro, hemp seed meal, soybean meal, and urea, beets yield with the application of the GSR G4 was higher by 685 lbs/acre, 1,656 lbs/acre, 1,453 lbs/acre, and 906 lbs/acre respectively. The beet yield with GSR G3 was higher by 373 lbs/acre compared to hemp seed meal, and 170 lbs/acre than soybean meal.

*Application rate: 30 lbs N ac<sup>-1</sup>*  
*Planting date: 12-Aug*  
*Harvest date: 12-Oct*  
*Crop: Beets (var Eagle)*  
*Soil Type: Benson rocky silt loam*  
*Row spacing: 40" between rows*



Field trials and crop growth. Image: Northwest Crops and Soils Program

There were no statistical differences in percent of organic matter and pH by treatment. Most soil nutrients declined from the beginning to the end of the season. The increase in pH from 6.93 on 14-Jun to 7.06 on 27-Oct was significant and likely attributed to the degradation of the organic matter over time. The decrease in organic matter from 4.73% to 4.38% was also significant and possibly due to the multiple tillage operations throughout the season. Additionally, phosphorous (P), potassium (K), sulfur (S), manganese (Mn), zinc (Zn), and boron (B) all had significant differences between the two sampling dates. All of the nutrient levels, except copper (Cu) which remained the same, decreased over the course of the season. This is most likely due to the crops using these nutrients and reducing their levels in the soil. Magnesium (Mg), calcium (Ca), sulfur (S), iron (Fe), and boron (B) differed statistically by treatment. Soybean meal had the highest concentration of Mg at 158 ppm, being statistically similar to GSR 3 and GSR 4. Soybean meal also had the highest concentration of Ca at 2744 ppm, being statistically similar to GSR 3, urea, and the control. Additionally, soybean meal had the highest concentration of B at 0.780 ppm, being statistically similar to GSR 3, GSR 4, and the control. Pro-Gro had the highest concentration of S at 10.4 ppm. Finally, GSR 3 had the highest Fe concentration of 3.94 ppm, being statistically similar to GSR 4, soybean meal, and the control. Additional field trials of GSR fertilizer varieties will be conducted.

Northwest Crops and Soils Program | 278 So. Main Street, Suite 2 | St. Albans, VT 05478

This work was supported by the United States department of Agriculture, Conservation Innovation Grant program, Natural Resources Conservation Service.

802-524-6501 or 1-800-639-2130 (toll-free in Vt.) | [susan.brouillette@uvm.edu](mailto:susan.brouillette@uvm.edu)

[uvm.edu/extension/nwcrops](http://uvm.edu/extension/nwcrops)