



United States
Department of
Agriculture

Forest Service

**Northern
Research Station**

General Technical
Report NRS-20



Ozone Bioindicator Sampling and Estimation

**Gretchen C. Smith
William D. Smith
John W. Coulston**



Abstract

Ozone is an important forest stressor that has been measured at known phytotoxic levels at forest locations across the United States. The percent forest exhibiting negative impacts from ozone air pollution is one of the Montreal Process indicators of forest health and vitality. The ozone bioindicator data of the U.S. Forest Service Forest Inventory and Analysis Program (FIA) are the only source of information available that documents plant injury from air pollution using consistent protocols. This document introduces the FIA ozone indicator and describes the sampling and estimation procedures of the national biomonitoring program. We provide background material on ozone, examples of bioindicator summary statistics, a description of spatial interpolation, and methods to estimate status and change in forested areas with respect to the occurrence of ozone injury from ambient ozone concentrations. The goal is to provide guidance to analysts and researchers on ways to incorporate ozone bioindicator data into reports and research studies. Periodic recommendations to analysts on improved analytical techniques will be made. Analysts are encouraged to consult the companion user guide for additional guidance on interpreting the ozone biomonitoring data and reporting on the issue of ozone and forest health for the FIA program.

Cover Photos

Front:

Jay Lackey, Forest Health Specialist at the Vermont Department of Forests and Parks, is shown evaluating bioindicator species for ozone injury in a typical open area suitable for ozone biomonitoring in eastern forest types.

Back:

Top left: Classic upper-leaf surface ozone stipple on black cherry. In the field, injury severity ranges from slight to severe on the same and different plants depending on genotype, leaf age, and ozone exposure.

Top right: Black cherry leaves showing obvious ozone injury and other symptoms. Three injured leaves are collected at random from individual trees of each injured species on each biomonitoring site.

Bottom: A plant press is used in the field to preserve the leaf samples with ozone injury. Pressed leaf samples are mailed to a regional ozone expert for microscopic examination and validation of the ozone symptom.

Published by:
USDA FOREST SERVICE
11 CAMPUS BLVD SUITE 200
NEWTOWN SQUARE PA 19073-3294

December 2007

For additional copies:
USDA Forest Service
Publications Distribution
359 Main Road
Delaware, OH 43015-8640
Fax: (740)368-0152

Visit our homepage at: <http://www.nrs.fs.fed.us/>

CONTENTS

Introduction.....	1
Overview of the Ozone Indicator	1
Detection Monitoring	2
Evaluation Monitoring	2
Key References	3
Biomonitoring and Data Collection	4
Point-In-Time Estimation	5
Plant-level Estimates	5
Biosite Index and Proportion Injured Plants	6
Status Estimation.....	6
Spatial Interpolation of the Biosite Index.....	7
Estimating Status for Forested Areas	8
Discussion	11
References	12
Appendices	17
Appendix I: Output Tables and Maps	18
Appendix II: Ozone Sensitivity of Tree and Shrub Species	24
Appendix III: Documents for Ozone Information Management	27

INTRODUCTION

Overview of the Ozone Indicator

Ozone interacts with forest ecosystems, causing visible injury and alterations in species composition and pest interactions (Chappelka and Samuelson 1998, Miller and others 1996). It is the only regional gaseous air pollutant that has been measured at known phytotoxic levels at both remote and urbanized forest locations (U.S. EPA 1996a,b). The importance of ozone as a forest stressor is illustrated by its inclusion in the Montréal Process Criteria and Indicators (Montreal Process 1995) in which the percent forest exhibiting negative impacts from air pollutants such as ozone is an indicator of the overall forest health and vitality. Coulston and others (2004) point out that the ozone biomonitoring data of the USDA Forest Service Forest Inventory and Analysis Program (FIA) are the only source of information available that documents plant injury from air pollution using consistent protocols. The goal of our document is to describe the ozone bioindicator and suggest analytical techniques appropriate for FIA ozone biomonitoring data.

The ozone bioindicator provides a biological index of ozone stress to plants using consistent protocols on a nationwide system of biomonitoring sites. Ozone biomonitoring is part of the FIA phase 3 sample (USDA Forest Service 2005) and is based on the documentation of visible foliar injury to known ozone-sensitive plant species under conditions of ambient exposure. The field methods, site variables, and site-level biosite index were developed with support from the scientific research community (Smith 1995), and the sampling procedures and analytical techniques have been reviewed in the scientific literature (Coulston and others 2003, Smith and others 2003).

Although ozone biomonitoring is part of the phase 3 (forest health) sample, the biomonitoring grid and the FIA base grid are independent and information is collected on a different population from other FIA phase 2 (P2) and phase 3 (P3) measurements (see Biomonitoring and Data Collection, page 4). One primary use of these data is Detection Monitoring (DM), which identifies forested areas that may be at risk from some stressor or disturbance (Detection Monitoring section, page 2). These areas may then become candidates for Evaluation Monitoring (EM) (see Evaluation Monitoring section, page 2). Vose (2000) defined risk analysis as the process of quantifying, either qualitatively or quantitatively, the probability and the potential impacts of some risk. DM analyses using the ozone biomonitoring data are both qualitative and quantitative and fall under the category of risk analysis.

Throughout this document the word “site” or “biosite” will be used to refer to the ground location where the FIA field crews collect ozone indicator data and the word “plot” will be used to refer to the FIA ground sample plots for which the estimation procedures apply. Our objective here is to provide guidance on presenting annual summaries of the ozone biomonitoring data and performing risk analysis for DM.

GRETCHEN C. SMITH,
*FIA National Ozone Advisor, University
of Massachusetts, Department of Natural
Resources Conservation, Amherst, MA.*
gcsmith@nrc.umass.edu

WILLIAM D. SMITH,
*Assessment Coordinator, U.S. Forest
Service, Southern Research Station,
Research Triangle Park, NC.*

JOHN W. COULSTON,
*Supervisory Research Forester, U.S. Forest
Service, Southern Research Station,
Knoxville, TN.*




Figure 1.—FIA ozone biomonitoring grid developed from the Environmental Monitoring and Assessment Program (EMAP) base grid (White and others 1992). The grid has four sampling intensities based on sensitive species and ambient ozone concentrations (see Smith and others 2001 for more details).

Detection Monitoring

The FIA ozone biomonitoring program is designed to detect and monitor plant-damaging concentrations of ozone in the natural environment. Information gathered at ozone biomonitoring sites identifies whether conditions exist (ozone, light, moisture, relative humidity) for plant injury to occur. This information can be used to report national and regional trends in ozone injury to plants and to identify areas of concern for closer evaluation. The biomonitoring grid is independent of the FIA base grid (Fig. 1). Therefore, spatial interpolation is one method to predict potential risk of ozone injury on the ground plots. Interpolated biosite values are classified into four response categories that are used to define and describe possible impact (i.e., risk) to the forest resource from ambient ozone exposure (Table 1). These categories also provide an indication of ozone relative air quality with respect to a plant receptor. The categorizations of the biosite index are qualitative, but have received positive reviews in the scientific literature (Coulston and others 2003, Smith and others 2003).

The categorized and interpolated biosite values, along with the presence and abundance of ozone-sensitive tree species found on FIA P2/P3 ground plots, are used to develop species-specific risk maps of ozone stress. These maps are then used to identify localized areas of moderate to high risk where EM studies are warranted. This approach is documented in Coulston and others (2003).

Evaluation Monitoring

When regional or national analyses of detection-level data indicate areas of potential impact on forest productivity and sustainability, these areas are evaluated through the implementation of additional studies on an intensified grid. For example, the biomonitoring data may indicate a band of high ozone stress across a state or region, which may prompt the regional analyst to ask if the finding is real or some artifact of data collection. The next logical step is an evaluation study to take a closer look at the area of concern. For the ozone indicator, an evaluation study should include an intensified

Table 1.—Classification scheme for the FIA biosite index¹

Biosite index	Bioindicator response	Assumption of risk	Possible impact	Relative air quality ²
0 - < 5	Little or no foliar injury	None	Visible injury to highly sensitive species, e.g., black cherry	Good
5 - < 15	Light to moderate foliar injury	Low	Visible injury to moderately sensitive species, e.g., tulip poplar	Moderate
15 - < 25	Moderate to severe foliar injury	Moderate	Visible and invisible injury. Tree-level response ³	Unhealthy for sensitive species
≥ 25	Severe foliar injury	High	Visible and invisible injury. Ecosystem-level response ³	Unhealthy

¹The categorizations of the biosite index are subjective and based solely on expert opinion.

²Relative ozone air quality from a plant's perspective. See: EPA-456/F-99-002 July 1999; <http://www.airnow.gov>.

³According to the EPA's Proposed Guidelines for Ecological Risk Assessment (Federal Register 61(175): 47552-47631).

sampling grid and an analysis of air quality and environmental data that influence plant response to ozone. It may also be important to examine species distribution maps available from FIA regional archives. An example of DM identifying a potential problem area and EM verifying the problem is illustrated by Coulston and others (2003) and Skelly and others (2003), respectively.

Key References

Biomonitoring: Biomonitoring has been used since the 1960s in numerous smaller field studies to assess pollutant stress. What is new and significant about the FIA biomonitoring program is its national scope and the successful implementation of national standards for training, field procedures, quality assurance, data analysis, and reporting.

Key references: Berry and Hepting 1964, Chappelka and others 1997, Feder 1978, Heck 1968, Hildebrand and others 1996, Kohut and others 1997, Lewis and Conkling 1994, Neufeld and others 1992, Pronos and Vogler 1981.

Bioindicator species: The FIA list of ozone bioindicator species was gleaned from various sources including the peer-reviewed scientific literature, interagency reports, and communications with Federal and university researchers experienced in ozone biomonitoring. Selected species are common across a variety of forest types, easy to identify and distinguish from similar species, and sensitive to ozone based on a combination of field evidence and causative fumigation experiments. The eastern bioindicator species have a long history of application in ozone field studies. The western bioindicator species have not been as well tested under natural conditions of ozone exposure, but have all received enough testing to justify inclusion in the FIA program.

Key references: East: Krupa and others 1998, Skelly and others 1987, Skelly 2000. West: Brace and others 1999, Campbell and others 2000, Duriscoe and Temple 1996, Mavity and others 1995, Temple 2000. East and West: U.S. Department of the Interior 2003.

Biosite index: The Horsfall-Barrett (HB) rating scale used to assess ozone injury in the field is based on a technique developed for plant disease research. Since the 1940s, it has been used repeatedly in the field evaluation of ozone-induced foliar injury. Details on the formulation of the site-level biosite index (BI) are presented in Biosite Index and Proportion Injured Plants, page 6. The index developed for the FIA program is new, but it has been widely adopted by cooperating researchers at various institutions and published in the scientific literature.

Key references: Horsfall and Barrett 1945, Horsfall and Cowling 1978, Smith and others 2003.

Classification scheme for the biosite index: The classification scheme for the FIA biosite index has been reviewed in the scientific literature and applied in a published assessment of ozone injury to eastern forest tree species.

Key references: Coulston and others 2003, Smith and others 2003.

Interpolation techniques: Plot-level attributes required for population estimates are developed by spatial interpolation of the biosite data. Spatial interpolation techniques are widely used in the analysis of air pollution, environmental, and ecological data. The approach used for the ozone indicator has been reviewed in the scientific literature and applied in a published assessment of ozone injury to eastern forest tree species.

Key references: Coulston and others 2003, Cressie 1993, Cressie and Ver Hoff 1993, Isaaks and Srivastava 1989, Lefohn and Pinkerton 1988.

Status and change estimation: There are several references for valid estimation techniques; however, Bechtold and Patterson (2005) provide a review and recommendations specifically for FIA data.

Key reference: Bechtold and Patterson (2005)

Biomonitoring and Data Collection

Ozone sampling occurs on a unique national grid independent of the P2 and P3 plot system. The ozone grid enhances quality assurance for this indicator by allowing greater flexibility in site location on the ground and greater sampling intensity in areas believed to be at high risk for ozone impact (Smith and others 2001). The grid design generates differing sampling intensities across the landscape based on the best available information on air quality regimes.

The ozone grid is purposive both at the grid level and at the biosite. Biosite location on the ground is deliberately chosen first for ease of access and second for optimal size, species, and plant counts. The ozone biomonitoring sites vary in size and do not have set boundaries. They are defined by the presence of ozone-sensitive bioindicator species indigenous to each FIA region. There must be one biosite per polygon on the national ozone biomonitoring grid. Some states use an intensified ozone grid so two or more biosites may be located in each polygon on the base grid. Biosite locations are mapped, geographic coordinates are recorded, and the same sites are evaluated every year. Ozone injury and our ability to detect that injury increase over the course of the field season. For this reason, the sampling window for the ozone indicator is limited to 3 weeks (from late-July to mid-August) within which the indicator is considered stable. This minimizes variability and the error associated with the data collection system.

The sampling rules for the ozone biosites are as follows. Biosites are wide-open areas, at least one acre in size, within or alongside forested areas. Each site must contain at least 30 individual plants of at least two bioindicator species. If not enough plants are available at one location, two nearby open areas, within 3 miles of each other, may be combined to maximize plant counts. Biosite locations must be easy to access, and they must be free of significant soil compaction and other human-made disturbance. Additional guidelines are available in the Field Methods Guide (USDA Forest Service 2000).

The characteristics of each site are described in terms of the size of the open area, elevation, terrain position, aspect, soil drainage, soil depth, and site disturbance. If characteristics vary significantly across the biosite, then the area where most of the bioindicator species are growing is described and variations are recorded on the site map and notes. When two nearby open areas are used, each location is described separately.

Up to 30 plants of each species are randomly selected for injury evaluation. Plants less than 12 inches in height, suppressed, shaded, or with more than half the crown out of reach are not evaluated. The approximate locations of the plants used for evaluations are drawn on the site map so that the same population of plants is evaluated on return visits to the biosite. The entire open area is sampled until 30 plants of two (ideally, three or more) species have been evaluated.

Quality assurance (QA) procedures dictate that the ozone injury symptom must be verified for each injured species on each site. Crews collect a minimum of three injured leaves from a random sample of individual plants that show obvious ozone injury, and they mail pressed leaf samples to a regional expert for review. Three leaves from each injured species are subject to microscopic examination. Injury is validated for all samples that show a characteristic color and injury pattern for ozone and that are otherwise free of confounding signs and symptoms of other mimicking stress agents (e.g., insects, disease, mites, or weather). If the symptoms are not typical of ozone injury, then the field data associated with the invalidated leaf voucher are zeroed out. Furthermore, if a leaf voucher is missing and unable to be validated, then the field data associated with the missing voucher are flagged so they cannot be used in data summaries or analyses.

The ozone indicator is included in the FIA National QA Plan (USDA Forest Service 2004). Just before the sampling window, ozone training and certification sessions are held in each region. A minimum of 10 biosites per region are blind checked every year (5 to 6 percent of the total biosites in each region). The ozone remeasurement data have been evaluated on two occasions, once in 1999 and again in 2003 (Pollard and Smith 2001, Pollard 2004). Inconsistent results with two eastern bioindicator species reported in Pollard and Smith (2001) were corrected by improvements to the ozone training session. Results from the 2003 review indicate the biosite data are robust and field crews in all regions are able to meet data quality expectations for the ozone indicator.

POINT-IN-TIME ESTIMATION

Plant-level Estimates

At each ozone biosite, 30 individual plants of two bioindicator species and between 10 and 30 individual plants of additional bioindicator species are evaluated for ozone injury. Each plant is rated for the proportion of leaves with ozone injury (injury amount) and the mean severity of symptoms (injury severity) using a modified Horsfall-Barratt scale with break points at 0, 6, 25, 50, 75, and 100

percent (Horsfall and Cowling 1978, USDA Forest Service 1999). This scale uses class break points that correspond to the ability of the human eye to distinguish gradations of healthy and unhealthy leaf tissue.

Biosite Index and Proportion Injured Plants

For each biosite, the percent injured plants and a biosite index are calculated based on the injury amount and severity scores. The proportion injured is $I_p = n_i/n_t$ where I_p is the proportion of plants injured, n_i is the number of injured plants (i.e., amount of injury $\neq 0$), n_t = the total number of plants evaluated. The biosite index is the average score (amount * severity) for each species averaged across all species on the biosite multiplied by 1,000 to allow risk categories to be defined by integers (Table 1). The biosite index is calculated

$$BI = 1000 \left(m^{-1} \sum_{j=1}^m n_j^{-1} \sum_{p=1}^{n_j \geq 10} a_{pj} s_{pj} \right)$$

where

BI = biosite index

m = number of species evaluated

n_j = number of plants of the j^{th} species evaluated

a_{pj} = proportion of injured leaves on the p^{th} plant of the j^{th} species

s_{pj} = average severity of injury on the p^{th} plant of the j^{th} species

Biosite summary statistics on the ozone indicator are generated annually and loaded to three ozone data summary tables in each FIA region. Tabular data include species and site counts and calculated mean injury indices from the first year each state implemented the ozone indicator up to the current year. For some regions it is important to group states with similar air quality regimes together and keep them separate from neighboring states with distinctly different air quality regimes. A map of biosite-level values will also be produced for illustration. Tables and figures illustrating these types of products are included in Appendix I.

Status Estimation

FIA plot-level attributes required for population estimates can be developed by spatial interpolation of data collected from the biosites. Each ozone season is unique, influenced by variable ozone levels, weather, windflow, and precipitation patterns. Therefore, it is important to use 5-year averages of the biosite index to generate a truly representative estimate of ozone stress. Thus, a 5-year moving average is used:

$$\hat{B}_t = \sum_{t=-(n-1)}^0 \frac{B_t}{n} \quad \text{where}$$

\hat{B}_t = the 5-year average estimate for a biosite value at time t

B_t = biosite value at time t

t = time in years ranging from $t = 0$ (the current year) to year $t - n$

n = number of biosite measurements (maximum of 5)

\hat{B}_t will be used in the spatial interpolation. Spatial interpolation techniques are widely used (e.g., see Cressie and Ver Hoff 1993) and air pollution variables are commonly interpolated. For example, Lefohn and Pinkerton (1988) interpolated ambient ozone concentrations to characterize forested

areas of the United States. Coulston and others (2003) interpolated data from ozone biomonitoring sites to characterize risk to northeastern tree species. For a review of spatial statistics, see Cressie (1993).

Spatial Interpolation of the Biosite Index

Many spatial interpolation techniques are available. In this document we discuss the procedures for kriging, inverse distance weighting, and cross-validation, but as work proceeds with the ozone indicator, other methods such as universal kriging, splining, and spatial regression may be implemented. Spatial interpolation is performed to create a map of ozone risk to plants. This map is used to, among other things, classify ozone injury risk for FIA P2/P3 plots. Many spatial interpolation techniques require analysts to make assumptions (e.g., stationarity). We assume analysts are aware of both the theoretical and practical considerations associated with each interpolation technique.

Using kriging, a standard interpolation technique, requires at least three steps to interpolate a surface. First, the empirical semivariogram is calculated. Second, the empirical semivariogram is modeled. With parameters from the modeled semivariogram, the kriging equations can be used. The semivariance between values for a particular lag distance h is $\gamma(h) = 1/(2N(h))\sum(v_i - v_j)^2$ where N is the number of pairs (i, j) and $v_i - v_j$ is the difference between the values of pair (i, j) . It is one-half the average squared difference between values a particular distance apart. Several model types can be used to model the empirical semivariogram. They include the Gaussian model, wave model, and exponential model. Gaussian models tend to account for strong spatial relationships at short distances while wave models account for periodicity in spatial relationships. Matern class models may also be used when flexibility near $h=0$ is desired. See Hoeting and others (2006) for more information on semivariogram model selection.

After the semivariogram has been modeled, ordinary kriging can be used to interpolate between values. Ordinary kriging is a weighted average such that $\hat{V}_0 = \sum_{i=1}^n w_i V_i$ where \hat{V}_0 is the estimate at unmeasured location 0 , w is the weight of each i observation, and V_i is the value of each i observation. The weights sum to 1 and are determined by minimizing the estimation error. The estimation variance is $S_0^2 = w_i \gamma(s_i - s_0) + \lambda$ where $\gamma(s_i - s_0)$ is the modeled semivariance for the distance between s_i and s_0 ; λ is the Lagrange multiplier from solving the linear system of equations for minimum estimation error.

Analysts may choose to use the inverse distance squared weighting interpolation method (IDW). Estimates are made by:

$$\hat{V}_0 = \frac{\sum_{i=1}^n d_{0i}^{-2} v_i}{\sum_{i=1}^n d_{0i}^{-2}}$$

where \hat{V}_0 is the estimate at unmeasured location 0 , d_{0i} is the distance from the i^{th} biosite to location 0 , and v_i is the value at biosite i . Under the assumption of intrinsic stationarity, the estimation variance is:

$$S_0^2 = -\sum_i \sum_j \left(\frac{d_{0i}^{-2}}{\sum d_{0i}^{-2}} \right) \left(\frac{d_{0j}^{-2}}{\sum d_{0j}^{-2}} \right) \gamma(s_i - s_j) + \sum_i \left(\frac{d_{0i}^{-2}}{\sum d_{0i}^{-2}} \right) 2\gamma(s_i - s_0).$$

Cross-validation is a method to quantify and compare various models (e.g., kriging and IDW). It can also be used to decide among variogram models (e.g., spherical, Gaussian). The cross-validation technique is implemented sequentially by removing each v_i one at a time and then estimating v_i based on the spatial model (e.g., IDW) and the remaining $n-1$ observations. If this is done sequentially for all $i = 1, \dots, n$ observations in the sample, the estimates can then be compared to the actual values using several standard summary statistics (Prediction error sum of squares – PRESS statistics).

The PRESS statistics are the values analysts may use to decide on which interpolation model performs the best for their particular situation. One PRESS statistic is the average squared deviation $= n^{-1} \sum_i (v_i - \hat{v}_{-i})^2$ where \hat{v}_{-i} is the prediction of v_i from the rest of the data. This value should be relatively small if the model fits well. Another summary statistic is the mean of standardized PRESS residuals $= n^{-1} \sum_i (v_i - \hat{v}_{-i}) / \sqrt{S_{(-i)}^2}$ where $S_{(-i)}^2$ is the estimation variance for \hat{v}_{-i} . This quantity should be close to zero if the model fits well. The root mean squared prediction residuals also provide a measure of model aptness. This is calculated by $\sqrt{n^{-1} \sum_i (v_i - \hat{v}_{-i} / \sqrt{S_{(-i)}^2})^2}$ and will be approximately one if the spatial model fits well. An analyst should create several interpolated maps using the various options for the IDW and kriging. For example, analysts may choose to create an IDW map based on the 12 nearest neighbors rather than all neighbors. Analysts may also decide to try several variogram models (e.g., spherical, Gaussian) with the kriging technique. The resultant maps can then be compared based on the PRESS statistics, and the analysts can decide on the most appropriate map.

Once an appropriate spatial model has been selected, biosite index values will be estimated for all P2 and P3 plots by intersecting the map of interpolated values with P2 and P3 plot locations (e.g., Fig. 2). This will result in a biosite index value estimate for each P2 and P3 plot (e.g., Table 2).

Table 2.—Example of interpolated biosite index values for P2 and P3 plots

Plot number	Biosite index	Injured plants (%)
27120110311029	13.8	15.8
27120110311156	16.1	18.0
27120110319064	0.1	1.1
27120110319251	20.8	40.8
27120110319361	19.9	25.0
27120110319385	9.7	10.4
27120110712093	1.7	3.8
27120110712438	8.4	8.0
27120110712720	6.1	7.9
27120110712907	24.5	30.8
27120110713096	10.6	10.0
27120110713107	20.9	14.5
27120110713459	14.0	12.9
27120110759099	13.6	6.8
27120110759237	2.8	5.0

Estimating Status for Forested Areas

Bioindicator attributes will be estimated yearly for all FIA plots, using the procedures described above. The attributes will then be merged with the other plot attributes. Population estimates include (1) the proportion of forest land in each biosite index category by region, ecoregion, and state; (2) the acres of forest land in each biosite index category by region, ecoregion, and state; and (3) the volume of ozone-susceptible species in each biosite index category by region, ecoregion, and state. Population estimates will be made using the procedures presented by Bechtold and Patterson (2005).




Figure 2.—Example of block kriging and intersecting Environmental Monitoring and Assessment Program (EMAP) hex centers.

The following description uses the same equations and terminology as used in “Sample-Based Estimators Used by the Forest Inventory and Analysis National Information System” (chapter 4 in Bechtold and Patterson 2005). In general terms, each P2 plot (and tree) will be assigned a biosite index category (Table 1) based on spatial interpolation that will be considered a plot (and tree) attribute. Each plot is assigned to one stratum from phase 1 (e.g., forest, nonforest). To estimate the proportion of forest land in biosite index category 4, an indicator function (δ) would be used. In this case, the indicator function would equal 1 if the attribute (biosite index category) is in the domain d (biosite index = 4) of interest, or 0 otherwise. The portion of each plot in the domain of interest is then

$$P_{hid} = \frac{\sum_j^4 \sum_k^{K_{hj}} a_{mhijk} \delta_{hijkd}}{a_m \bar{P}_{mh}} \text{ where}$$

P_{hid} = proportion of plot i in the domain of interest d , for plots assigned to stratum h , adjusted for stratum h plots that overlap the population boundary

a_{mbijk} = mapped area (acres) of subplot (macroplot) j covering condition k on plot i assigned to stratum h . (Area is computed using the largest area mapped, which is the subplot except in the Pacific Northwest (PNW) where the macroplot or 1-ha circle is used.)

δ_{hijkd} = zero-one domain indicator function, which is 1 if condition k on subplot (macroplot) j of plot i assigned to stratum h belongs to the domain of interest d

K_{hij} = the number of conditions that exist on subplot (macroplot) j of plot i assigned to stratum h

a_m = total area of the largest plot on which area attributes are mapped (i.e., four times the subplot or macroplot area)

\bar{p}_{mh} = mean proportion of stratum h mapped plot areas falling within the population (\bar{p}_{mh} is generally 1 unless the plot is partially outside the population. If this situation arises, see Bechtold and Patterson (2005)).

The estimated proportion of forest land in strata h and domain d is simply the average of the plot values.

$$\bar{P}_{hd} = \frac{\sum_i^{n_h} P_{hid}}{n_h}$$

The total area in the domain of interest is then

$$\hat{A}_d = A_T \sum_h^H W_h \bar{P}_{hd} = A_T \bar{P}_d \quad \text{where}$$

A_T = total area in the population in acres

\bar{P}_d = estimated proportion of the population in the domain of interest d

W_h = weight for stratum h , that is, the proportion of the population area, A_T , that is in stratum h

A similar procedure is used to estimate the volume of susceptible tree species in each ozone biosite index category. Tables 1 and 2 in the section Ozone Sensitivity of Tree and Shrub Species provide a preliminary list of tree and shrub species susceptible to ozone injury. As an example, suppose the attribute of interest was the total volume of loblolly pine in biosite index category 4. The following is used to estimate the attribute of interest on a per unit area basis:

$$y_{hid} = \frac{\sum_j^4 \sum_t y_{hijt} \delta_{hijtd}}{a_o \bar{p}_{oh}} \quad \text{where}$$

y_{hijt} = attribute of interest for tree t on macroplot, subplot or microplot j of plot i assigned to stratum h

δ_{hijtd} = zero-one domain indicator function, which is 1 if tree t on subplot j of plot i assigned to stratum h belongs to the domain of interest d

a_o = total area normally used to observe the attribute of interest on a plot, that is, four times the microplot, subplot, or macroplot area

$$\bar{p}_{oh} = \frac{\sum_i^{n_h} \sum_j^4 \sum_k^{K_{hij}} a_{ohijk} \delta_{hijk}}{a_o n_h} = \frac{1}{a_o n_h} \sum_i^{n_h} \sum_j^4 \sum_k^{K_{hij}} a_{ohijk} \delta_{hijk} = \text{mean proportion of stratum } h \text{ observed-plot areas}$$

falling within the population where, a_{ohijk} = area normally used to observe the attribute of interest (microplot, subplot or macroplot j) covering condition k on plot i assigned to stratum h .

These values are then averaged across each i plot

$$\overline{Y}_{hd} = \frac{\sum_i^{n_h} y_{hid}}{n_h}$$

The total for the attribute of interest in the domain of interest is then

$$\hat{Y}_d = A_r \sum_h^H W_h \overline{Y}_{hd} = A_r \overline{Y}_d$$

See Output Tables and Maps section for example output.

DISCUSSION

Here we present one method to perform DM by classifying each FIA plot based on an interpolated map of ozone injury risk. The purpose of this activity is to identify candidate areas for EM. As with other DM activities, there is a high noise to signal ratio and there may be a relatively high rate of false positives. For this reason, EM is an essential part of the process. The map of ozone injury risk does have unquantified error. However, other maps used to classify FIA plots (e.g., ecoregion sections, counties) also have unquantified error. When the information is used at its intended resolution, unquantified errors may be overlooked. For the ozone bioindicator, error propagation can be overlooked for DM activities. However, error propagation cannot be overlooked if one is trying to make a statistical inference about the relationship between growth rates and ozone injury.

The purpose of this document is to describe the analytical techniques used with the ozone indicator. We provided background material on ozone, examples of biosite summary statistics, a description of spatial interpolation, and methods to estimate status and change in forested areas with respect to the occurrence of ozone injury from ambient ozone concentrations. Appendix I includes examples of each expected output. Appendix II provides a current list of tree and shrub species susceptible to ozone. Appendix III contains several additional documents to assist FIA analysts with ozone data access and management. The list in Appendix II will be updated as more information becomes available. The interpolation techniques may be improved over time, and other methods of estimating change (e.g., spatio-temporal kriging) may also be investigated. Periodic recommendations to analysts will be made as results for QA analyses become available. There is also a companion ozone bioindicator user guide (Smith and others, in review) that analysts are encouraged to consult for additional guidance on interpreting ozone biomonitoring data and reporting on the issue of ozone and forest health for the FIA program.

ACKNOWLEDGMENTS

The ozone biomonitoring program began in the early 1990s and has succeeded due to the efforts and direction of numerous investigators, regional trainers, and state forest health coordinators. The authors wish to extend special thanks to the following individuals: Ed Jepsen, WI Department of Natural Resources; Dan Stratton and John Simpson, U.S. Forest Service, Southern Research Station; Barbara O'Connell, U.S. Forest Service, Northern Research Station; Sally Campbell, U.S. Forest Service, Pacific Northwest Research Station; and Patrick Temple, Ecological Systems Group, Riverside, CA. We also thank the regional field crews and quality assurance staff for their efforts and commitment to data quality. Without them the national ozone biomonitoring program would not be possible.

REFERENCES

- Bechtold, W.A.; Patterson, P.L., eds. 2005. **The enhanced forest inventory and analysis program-national sampling design and estimation procedures**. Gen. Tech. Rep. SRS-80. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 85 p.
- Berry, C.R.; Hepting, G.H. 1964. **Injury to eastern white pine by unidentified atmospheric constituents**. Forest Science. 10: 2-13.
- Brace, S.; Peterson, D.L.; Horner, D. 1999. **A guide to ozone injury in vascular plants of the Pacific Northwest**. Gen. Tech. Rep. PNW-446. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 63 p.
- Campbell, S.; Smith, G.; Temple P.; Pronos, J.; Rochefort, R.; Anderson, C. 2000. **Monitoring for ozone injury in west coast (Oregon, Washington, California) forests in 1998**. Gen. Tech. Rep. PNW-495. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 19 p.
- Chappelka A.; Renfro, J.R.; Somers, G.L. 1997. **Evaluation of ozone injury on foliage of black cherry (*Prunus serotina*) and tall milkweed (*Asclepias exalta*) in Great Smoky Mountains National Park**. Environmental Pollution. 95: 13-18.
- Chappelka, A.H.; Samuelson, L.J. 1998. **Ambient ozone effects on forest trees of the eastern United States: a review**. New Phytologist. 139: 91-108.
- Coulston, J.W.; Riitters, K.H.; Smith, G.C. 2004. **A preliminary assessment of Montreal Process Indicators of air pollution for the United States**. Environmental Monitoring and Assessment. 95: 57-74.
- Coulston, J.W.; Smith, G.C.; Smith, W.D. 2003. **Regional assessment of ozone sensitive tree species using bioindicator plants**. Environmental Monitoring and Assessment. 83: 113-127.
- Cressie, N.A.C. 1993. **Statistics for spatial data**. New York: John Wiley and Sons. 900 p.
- Cressie, N.A.C.; Ver Hoff, J.M. 1993. **Spatial statistical analysis of environmental and ecological data**. In: Goodchild, M.F.; Parks, B.O.; Steyaert, L.T., eds. Environmental modeling with GIS. New York: Oxford University Press. 488 p.
- Duriscoe, D.M.; Temple, P.J. 1996. **Ozone West Bioindicator Survey - results of a preliminary survey of some ozone sensitive plants in the West**. Unpublished report. On file with: USDA Forest Service, Forest Health Monitoring Program, Research Triangle Park, NC 27709
- Eckert, R.T.; Kohut, R.; Lee, T.; Staplefeldt, K. 1999. **Foliar ozone injury on native vegetation at Acadia National Park: results from a six-year (1992-1997) field survey**. Denver, CO: U.S. Department of the Interior, National Park Service. Air Resources Division. 42 p.

- Feder, W.A. 1978. **Plants as bioassay systems for monitoring atmospheric pollutants.** Environmental Health Perspectives. 27: 139-147.
- Gillespie, A.J.R. 1999. **Rationale for a national annual inventory program.** Journal of Forestry. 97: 16-20.
- Heck, W.W. 1968. **Factors influencing expression of oxidant damage to plants.** Annual Review of Phytopathology. 6: 165-187.
- Hildebrand, E.S.; Skelly, J.M.; Federickson, T. 1996. **Incidence of ozone-induced foliar injury on sensitive hardwood tree species from 1991-1993 in the Shenandoah National Park, Virginia.** Canadian Journal of Forest Research. 26: 658-659.
- Hoeting, J.A.; Davis, R.A.; Merton, A.A.; Thompson, S.E. 2006. **Model selection for geostatistical models.** Ecological Applications. 16(1): 87-98.
- Horsfall, J.G.; Barrett, R.W. 1945. **An improved grading system for measuring plant disease.** Phytopathology. 35: 655.
- Horsfall, J.G.; Cowling, E.B., eds. 1978. **Pathometry: the measurement of plant disease.** In: Plant disease. New York: Academic Press: 119-136. Vol. II.
- Isaaks, E.H.; Srivastava, R.M. 1989. **An introduction to applied geostatistics.** New York: Oxford University Press. 561 p.
- Kohut, R.; Laurence, J.; King, P.; Raba, R. 1997. **Identification of bioindicator species for ozone and assessment of the responses to ozone of native vegetation at Acadia National Park.** On file with: U.S. Department of the Interior, National Park Service, P.O. Box 25287, Denver, CO 80225-0287.
- Krupa, S.V.; Manning, W.J. 1988. **Atmospheric ozone: formation and effects on vegetation.** Environmental Pollution. 50: 01-137.
- Krupa, S.V.; Tonneijck, A.E.G.; Manning, W.J. 1998. **Ozone.** In: Flager, R.B. ed. Recognition of air pollution injury to vegetation: a pictorial atlas. Pittsburgh, PA: Air & Waste Management Association: sec. 2.1-2.23.
- Lefohn, A.S.; Pinkerton, J.E. 1988. **High resolution characterization of ozone data for sites located in forested areas of the United States.** JAPCA. 38: 1504-1511.
- Lewis, T.E.; Conkling B.L., eds. 1994. **Forest Health Monitoring southeast loblolly/shortleaf pine demonstration interim report.** Washington, DC: U.S. Environmental Protection Agency: sec. 3.1-3.8.

- Mavity, E.; Stratton, D.; Berrang, P. 1995. **Effects of ozone on several species of plants which are native to the western United States.** Unpublished report on file with: USDA Forest Service, Center for Forest Environment Studies, Dry Branch, GA. 12 p.
- Miller, P.R.; Stolte, K.W.; Duriscoe, D.M.; Pronos, J. 1996. **Extent of ozone injury to trees in the western United States.** In: Evaluating ozone air pollution effects on pines in the western United States. Gen. Tech. Rep. PSW-155. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 1-6.
- Montreal Process. 1995. <http://www.mpci.org>.
- Neufeld, H.S.; Renfro, J.R.; Hacker, W.D.; Silsbee, D. 1992. **Ozone in Great Smoky Mountains National Park: dynamics and effects on plants.** Proc. 85th annual meeting, Air & Waste Management Association; 1992 November 4-7; Atlanta, GA: TR-20 Tropospheric ozone and the environment II, effects, modeling, and control: 594-617.
- Pollard, J.E. 2004. **Analysis of data quality for the ozone indicator, biosite plot index and number of species observed by crews and auditor.** U.S. Department of Agriculture, Forest Service. On file with: Harry Reid Center for Environmental Studies, University of Nevada - Las Vegas, Las Vegas, NV 89154.
- Pollard, J.E.; Smith, W.D. 2001. **Forest Health Monitoring 1999 plot component quality assurance report.** On file with: USDA Forest Service, National Forest Health Monitoring Program, Research Triangle Park, NC 27709.
- Pronos, J.; Vogler D.R. 1981. **Assessment of ozone injury to pines in the southern Sierra Nevada, 1979/1980.** FPM Rep. 81-20. Berkeley, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Region. 13 p.
- Renfro, J.R. **Great Smoky Mountains National Park ozone fumigation program plant testing and sensitivity list from 1987-1992.** Unpublished report, available from: U.S. Department of the Interior, National Park Service, Upland Fields Research Laboratory, GSMNP, Gatlinburg, TN 37738.
- Skelly, J.M. 2000. **Tropospheric ozone and its importance to forests and natural plant communities of the northeastern United States.** Northeastern Naturalist. 7: 221-236.
- Skelly, J.M.; Davis, D.D.; Merrill W.; Cameron, E.A.; Brown, D.B.; Dochinger, L.S. 1987. **Diagnosing injury to eastern forest trees - A manual for identifying damage caused by air pollution, pathogens, insects, and abiotic stresses.** NAPAP Forest Response Program, National Vegetation Survey. Available from: Publications Distribution Center, The Pennsylvania State University, 112 Agricultural Administration Building, University Park, PA 16802.
- Skelly, J.M.; Yuska, D.J.; Savage, J.E.; Ferdinand, J.A.; Orendovici, F.; Stevenson, R. 2003. **An FHM evaluation monitoring project: investigation of factors associated with ozone-induced**

- foliar injury within biomonitoring plots in southwestern Pennsylvania forests.** PSIE 2003-5. University Park, PA: Pennsylvania State University Institute of the Environment.
- Smith, G.C. 1995. **FHM second ozone bioindicator workshop - summary of proceedings.** Unpublished report. On file with: USDA Forest Service, Forest Health Monitoring Program, Research Triangle Park, NC 27709.
- Smith, G.C.; Coulston, J.W.; Jepsen, E.; Prichard, T. 2003. **A national ozone biomonitoring program – results from field surveys of ozone sensitive plants in northeastern forests (1994 - 2000).** Environmental Monitoring and Assessment. 87: 271-291.
- Smith, G.C.; Coulston, J.W.; O'Connell, B. In review. **Ozone bioindicators and forest health: a guide to the evaluation, analysis, and interpretation of ozone injury data in the Forest Inventory and Analysis Program.**
- Smith, W.D.; Coulston, J.W.; Smith, G.C. 2001. **Development of the ozone indicator sampling grid.** Unpublished report. On file with: USDA Forest Service, Forest Inventory and Analysis Program, Research Triangle Park, NC 27709.
- Smith, W.H. 1981. **Forest stress: symptomatic foliar damage caused by air contaminants.** In: DeSanto, R.S., ed. Air pollution and forests, interactions between air contaminants and forest ecosystems. New York: Springer-Verlag. 379 p.
- Temple, P.J. 2000. **Ozone responses of bioindicator plants for forests in the western United States, Final report.** On file with: USDA Forest Service, Forest Health Monitoring Program, Research Triangle Park, NC 27709.
- USDA Forest Service. 1999. **Forest Health Monitoring 1999 Field Methods Guide.** USDA Forest Service, National Forest Health Monitoring Program. Research Triangle Park, NC 27709.
- USDA Forest Service. 2000. **Forest Inventory and Analysis, National Core Field Guide: Field data collection procedures for Phase 2 & 3 Plots – Ozone bioindicator plants.** U.S. Department of Agriculture, Forest Service, Northeastern Research Station, Newtown Square, PA, and Pacific Northwest Research Station, Portland, OR. Vol. 1, Sec. 9.0.
- USDA Forest Service. 2004. **Forest Inventory and Analysis, Plot Component Quality Assurance Implementation Plan, Version 1.0.** Internal report. On file with: U.S. Department of Agriculture, Forest Service, 201 14th St., Washington, DC 20250.
- USDA Forest Service. 2005. **Phase 2 and Phase 3: Ground Measurements.** FIA Fact Sheet Series. http://fia.fs.fed.us/library/fact-sheets/data-collections/Phase2_3.pdf.
- U.S. Department of the Interior. National Park Service Air Resources Division. U.S. Fish and Wildlife Service Air Quality Branch. Denver, CO. November 2003. NPS D1522. Natural Resources Report NPS/NRARD/NRR-2003/01. **Ozone sensitive plant species on National**

Park Service and U.S. Fish and Wildlife Lands: results of a June 24-25, 2003 workshop.
Baltimore, MD. Available at: <http://www2.nature.nps.gov/air/studies/ecoOzone.cfm>.

U.S. Environmental Protection Agency. 1996a. **Air quality criteria for ozone and related photochemical oxidants.** Environmental concentrations, patterns, and exposure estimates. EPA/600/P-93/004aF. Washington, DC: Office of Research and Development. Vol.1, Sec. 4.

U.S. Environmental Protection Agency. 1996b. **Air quality criteria for ozone and related photochemical oxidants.** Environmental effects of ozone and related photochemical oxidants, EPA/600/P-93/004aF. Washington, DC: Office of Research and Development. Vol. II, Sec .5.

Vose, D. 2000. **Risk analysis: a quantitative guide.** New York: John Wiley and Sons. 418 p.

White, D.; Kimerling, A.J.; Overton, W.S. 1992. **Cartographic and geometric component of a global sampling design for environmental monitoring.** Cartography and Geographic Information Systems. 19: 5-22.

APPENDICES

The appendices include supplementary information on the ozone indicator. Appendix I, Output Tables and Maps, includes examples of output tables and maps suitable for FIA state reports. Appendix II, Ozone Sensitivity of Tree and Shrub Species, includes ozone sensitivity tables for trees and shrubs, information that is needed for risk assessment analysis, and Appendix III, Documents for Ozone Information Management, provides information on ozone data in the FIA national information management system (NIMS) and FIA public database (FIADB), as well as contact information for individuals most familiar with the ozone biomonitoring program in FIA.

Appendix I: Examples of Output Tables and Maps for Annual and Multiyear Summary Reports

Table 3. State-level summary statistics

Table 4. Region-level summary statistics

Table 5. Example of summary statistics using real data

Table 6. County-level population estimates

Figure 3. National map of ozone risk to plants.

Figure 4. Example of State-level population estimates using real data.

Table 3.—State-level summary statistics

Parameter	State X -Biomonitoring Program								
	1994	1995	1996	1997	1998	1999	2000	2001	2002
Number of biosites evaluated	XX	XX	XX	XX	XX	XX	XX	XX	XX
Number of biosites with injury	X	XX	X	X	XX	X	X	XX	XX
Average biosite injury score ¹	X.X	X.X	X.X	X.X	X.X	X.X	X.X	X.X	X.X
Percent biosites with BI = 0 to 4.9 ²	XX	XX	XX	XX	XX	XX	XX	XX	XX
Percent biosites with BI = 5 to 14.9	XX	XX	XX	XX	XX	XX	XX	XX	XX
Percent biosites with BI = 15 to 24.9	XX	XX	XX	XX	XX	XX	XX	XX	XX
Percent biosites with BI >= 25	XX	XX	XX	XX	XX	XX	XX	XX	XX
Average number of species per biosite	X	X	X	X	X	X	X	X	X
Number of plants evaluated	XXXX	XXXX	XXXX	XXXX	XXXX	XXXX	XXXX	XXXX	XXX
Number of plants injured	X	XX	XX	XX	XXX	X	X	XXX	X
Percent sample plants by HB category ³									
0 = no injury	XX	XX	XX	XX	XX	XX	XX	XX	XX
1 = 1 to 6%	XX	XX	XX	XX	XX	XX	XX	XX	XX
2 = 7 to 25%	X	X	X	X	X	X	X	X	X
3 = 26 to 50%	X	X	X	X	X	-	X	X	X
4 = 51 to 75%	-	X	-	X	X	-	X	X	-
5 = >75%	-	X	-	X	-	-	X	X	-
Number of plants evaluated by species									
Species 1 (#injured in parentheses)	X (X)	X(X)	X(X)	X(X)	X(X)	-	X(X)	X(X)	X(X)
Species 2, etc.	-	X	X(X)	X(X)	X(X)	X	XX	X(X)	X

¹The biosite index is based on the average injury score (amount*severity) for each species averaged across all species on the biosite.

²Biosite categories represent a relative measure of tree-level response to ambient ozone exposure (see table 1 in the main body of the text).

³HB = injury severity is an estimate of the mean severity of symptoms on injured foliage (0 = no injury; 1=1-6%; 2 = 7- 25%; 3 = 26-50%; 4 = 51-75%; 5 >75%). Calculated percents are rounded to the nearest whole number. Terms are further described in the text.

*Standard errors can be presented, as needed, for the calculated variables.

Table 4.—Region-level summary statistics

Parameter	ABC Region - Biomonitoring Program								
	1994	1995	1996	1997	1998	1999	2000	2001	2002
Number of biosites evaluated	xx	xx	xx	xx	xx	xx	xx	xx	xx
Number of biosites with injury	x	xx	x	x	xx	x	x	xx	xx
Number of plants evaluated	xxxx	xxxx	xxxx	xxxx	xxxx	xxxx	xxxx	xxxx	xxx
Number of plants injured	x	xx	xx	xx	xxx	x	x	xxx	x
Average biosite injury score ¹	x.x	x.x	x.x	x.x	x.x	x.x	x.x	x.x	x.x
Percent biosites with BI = 0 to 4.9 ²	xx	xx	xx	xx	xx	xx	xx	xx	xx
Percent biosites with BI = 5 to 14.9	xx	xx	xx	xx	xx	xx	xx	xx	xx
Percent biosites with BI = 15 to 24.9	xx	xx	xx	xx	xx	xx	xx	xx	xx
Percent biosites with BI >= 25	xx	xx	xx	xx	xx	xx	xx	xx	xx
Average number of species per biosite	x	x	x	x	x	x	x	x	x
Number of plants evaluated	xxxx	xxxx	xxxx	xxxx	xxxx	xxxx	xxxx	xxxx	xxx
Number of plants injured	x	xx	xx	xx	xxx	x	x	xxx	x
Percent sample plants by HB category ³									
0 = no injury	xx	xx	xx	xx	xx	xx	xx	xx	xx
1 = 1 to 6%	xx	xx	xx	xx	xx	xx	xx	xx	xx
2 = 7 to 25%	x	x	x	x	x	x	x	x	x
3 = 26 to 50%	x	x	x	x	x	-	x	x	x
4 = 51 to 75%	-	x	-	x	x	-	x	x	-
5 = >75%	-	x	-	x	-	-	x	x	-
Number of plants evaluated by species									
Species1 (#injured in parentheses)	x (x)	x(x)	x(x)	x(x)	x(x)	-	x(x)	x(x)	x(x)
Species2,etc.	-	x	x(x)	x(x)	x(x)	x	xx	x(x)	x

¹The biosite index is based on the average injury score (amount*severity) for each species averaged across all species on the biosite.

²Biosite categories represent a relative measure of tree-level response to ambient ozone exposure (see table 1 in the main body of the text).

³HB = injury severity is an estimate of the mean severity of symptoms on injured foliage (0 = no injury; 1=1-6%; 2 = 7- 25%; 3 = 26-50%; 4 = 51-75%; 5 >75%). Calculated percents are rounded to the nearest whole number. Terms are further described in the text.

*Standard errors can be presented, as needed, for the calculated variables.

Note: Tables 3 and 4 provide an example of site-level summary statistics from State X and Region ABC. These two tables are core products for the ozone indicator. The summarized values show the base data used to generate the plot-level and population-level estimates as described in the text of this document. Individual states may choose to use the regional table as a basis of comparison to their summary statistics. Smaller states may choose to use the regional table for reports.

Table 5.—Number of biomonitoring sites evaluated for ozone-induced foliar symptoms, number of plants sampled, and percent of sampled plants in each injury severity category by year and subregion in FIA-North

Subregion and year ¹	No. of biosites evaluated	No. of plants sampled	Injury severity categories ²					
			0	1	2	3	4	5
Percent of sampled plants								
Lake States								
1996	95	3,880	99	1	<1	<1	0	0
1997	104	4,584	99	1	<1	<1	0	<1
1998	160	9,012	97	2	1	<1	<1	<1
1999	143	10,949	97	2	1	<1	<1	<1
2000	160	12,647	97	2	1	<1	<1	<1
New England								
1996	92	4,245	89	5	4	2	<1	<1
1997	91	4,248	93	3	3	1	<1	<1
1998	98	5,460	90	4	4	2	<1	<1
1999	96	5,057	97	1	1	<1	<1	<1
2000	87	4,850	96	2	2	<1	<1	0
North Central								
1996	8	589	67	6	7	4	7	9
1997	19	1,180	77	3	9	6	4	<1
1998	36	1,580	72	5	10	9	4	<1
1999	45	3,387	90	4	3	2	1	<1
2000	131	8,688	92	4	3	1	<1	<1
Mid-Atlantic								
1996	34	1,244	82	5	5	5	2	1
1997	60	2,908	93	2	2	2	1	<1
1998	170	6,384	78	5	7	5	3	2
1999	191	10,941	97	1	1	1	<1	<1
2000	182	12,762	93	2	2	1	1	<1

¹ Subregions are defined as follows: Lake States = MI, MN, WI; New England = CT, ME, MA, NH, RI, VT; North Central = IL, IN, IA, MO; Mid-Atlantic = DE, MD, NJ, OH, PA, WV.

² Injury severity is an estimate of the mean severity of symptoms on injured foliage (0 = no injury; 1=1-6%; 2 = 7- 25%; 3 = 26-50%; 4 = 51-75%; 5 >75%). Calculated percents are rounded to the nearest whole number.

Note: Table 5 is an example of site-level summary statistics using real data. It is sometimes useful to summarize ozone bioindicator data by multi-state groupings. For example, in the Northeast, it is informative to separate New England and New York from the Mid-Atlantic States because they tend to have dramatically different air quality regimes. Reporting a single regionwide injury index may mask gradations in air quality across the landscape and make it difficult to assess changes in the ozone indicator over time.

Table 6.—Population estimates for the ozone indicator including acres of forest land and volume of ozone-susceptible tree species in each biosite index category by state and county in Region ABC. Real data for Delaware (2002) are presented.

State	Cnty	Biosite Index ¹							
		Acres of forest land				Volume of susceptible species			
		<5	5 - <15	15 - <25	≥25	<5	5 - <15	15 - <25	≥25
10	1	0	97,356	56,338	11,119	xxx	xxx	xxx	xxx
10	3	0	3,954	71,164	69,434	xxx	xxx	xxx	xxx
10	5	0	7,166	97,356	35,088	xxx	xxx	xxx	xxx
State 2	1	xxx	xxx	xxx	xxx	xxx	xxx	xxx	xxx
State 2	2	xxx	xxx	xxx	xxx	xxx	xxx	xxx	xxx
State 2	3	xxx	xxx	xxx	xxx	xxx	xxx	xxx	xxx
State 3	1	xxx	xxx	xxx	xxx	xxx	xxx	xxx	xxx
State 3	2	xxx	xxx	xxx	xxx	xxx	xxx	xxx	xxx
State 3	3	xxx	xxx	xxx	xxx	xxx	xxx	xxx	xxx

¹The biosite index is based on interpolated values (see Plant-level Estimates, page 5, for details). Biosite categories represent a relative measure of tree-level response to ambient ozone exposure (see Table 1).




Figure 3.—National map of ozone risk to plants. Categorized values are derived from the 1999-2002 biosite data.

Note: This map of ozone risk to plants is a core product for the ozone indicator. In this example, biosite index values were averaged across the 4-year sampling period from 1999 to 2002 and then geostatistical procedures were used to create an interpolated bioindicator response surface across the landscape (see Status Estimation, page 6). The interpolated data are classified into color-based gradations of response representing low risk of probable ozone injury to forests (green), moderate risk (yellow), and high risk (red). These categories also provide an indication of ozone relative air quality with respect to a plant receptor (see Table 1). Intensified sampling is recommended where high ozone stress coincides with the spatial distribution of ozone-sensitive tree species. Refer to Appendix II for more information on the ozone sensitivity of tree species.

The ozone risk map is used to estimate bioindicator attributes for all FIA plots using the procedures described in the section on Spatial Interpolation of the Bioside Index, page 7. BI attributes are merged with other FIA plot attributes to generate population estimates using the procedures described in the section on Estimating Status for Forested Areas, page 8.

Biosite Index and Risk Estimation SOUTH CAROLINA




Figure 4.—Total tree volume, tree volume for ozone-sensitive trees, and sensitive volume as a percent of total volume by ozone risk category for South Carolina (2002).

Note: The data presented in Figure 4 provide an example of ozone risk estimation at the State level. Biosite categories on the x axis represent the risk of probable ozone injury to ozone-sensitive tree species in South Carolina in 2002. More than 16 million cubic feet of tree volume was categorized for risk of ozone injury. Approximately 6 million cubic feet falls into the no risk category while just over 10 million cubic feet of tree volume falls into the low to moderate risk categories. Fifty-three percent of the total categorized tree volume includes tree species that are ozone sensitive.

In this example, estimates are presented in terms of tree volume. However, other useful population estimates include the proportion of forest land and the acres of forest land in each biosite index category. Refer to Status Estimation, page 6 for the procedures used to estimate bioindicator attributes for forested areas.

Appendix II: Ozone Sensitivity of Tree and Shrub Species

The abbreviations used to assign sensitivity in the following tables are as follows: Sen = ozone sensitive, ModSen = moderately sensitive, InSen = ozone insensitive, Unk = unknown ozone sensitivity because there is evidence from different observers that is conflicting. Regional analysts should review both tables because species listed as eastern may be found in limited areas in Western States and visa versa. Additional ozone sensitivity listings of non-woody, forest species can be found at: <http://www2.nature.nps.gov/air/Pubs/pdf/BaltFinalReport1.pdf>.

Table 7. List of eastern tree and shrub species and their ozone sensitivity.

Table 8. List of western tree and shrub species and their ozone sensitivity.

Table 7.—List of eastern tree and shrub species and their ozone sensitivity

Eastern species		Sensitivity	Citation
balsam fir	<i>Abies balsamea</i>	InSen ¹	Smith 1981
boxelder	<i>Acer negundo</i>	ModSen ¹	Smith 1981
striped maple	<i>Acer pensylvanicum</i>	Unk	
red maple	<i>Acer rubrum</i>	Sen	Eckert et al. 1999
silver maple	<i>Acer saccharinum</i>	Unk	USDI 2003
sugar maple	<i>Acer saccharum</i>	InSen	Renfro 1987-1992
mountain maple	<i>Acer spicatum</i>	Unk	
Ohio buckeye	<i>Aesculus glabra</i>	Unk	USDI 2003
yellow buckeye	<i>Aesculus octandra</i>	Sen ²	USDI 2003
tree-of-heaven	<i>Ailanthus altissima</i>	Sen ²	USDI 2003
speckled alder	<i>Alnus rugosa</i>	Sen ²	USDI 2003
serviceberry	<i>Amelanchier arborea</i>	Sen	Renfro 1987-1992
Allegheny seviceberry	<i>Amelanchierlaevis</i>	Unk	USDI 2003
pawpaw	<i>Asimina triloba</i>	Unk	
yellow birch	<i>Betula alleghaniensis</i>	Sen	Renfro 1987-1992
sweet birch	<i>Betula lenta</i>	Unk	
paper birch	<i>Betula papyifera</i>	ModSen	Eckert et al. 1999
gray birch	<i>Betula populifolia</i>	ModSen	Eckert et al. 1999
bitternut hickory	<i>Carya cordiformis</i>	Unk	
pignut hickory	<i>Carya glabra</i>	Unk	
shagbark hickory	<i>Carya ovata</i>	Unk	
hickory sp.	<i>Carya sp.</i>	Unk	
mockernut hickory	<i>Carya tomentosa</i>	Unk	
hackberry	<i>Celtis occidentalis</i>	Unk	
common buttonbush	<i>Cephalanthus occidentalis</i>	Unk	USDI 2003
eastern redbud	<i>Cercis canadensis</i>	ModSen, Sen ²	Renfro 1987-1992, USDI 2003
yellowwood	<i>Cladrastis lutea</i>	Unk	USDI 2003
Virgin's bower	<i>Clematis virginiana</i>	Sen ²	USDI 2003
flowering dogwood	<i>Cornus florida</i>	ModSen	Renfro 1987-1992
American hazelnut	<i>Corylus americana</i>	Sen ²	USDI 2003
Hawthorn	<i>Crataegus sp.</i>	Sen ³	Krupa et al. 1998
common persimmon	<i>Diospyros virginiana</i>	Unk	
American beech	<i>Fagus grandifolia</i>	Unk	
white ash	<i>Fraxinus americana</i>	Sen	Skelly 2000
black ash	<i>Fraxinus nigra</i>	Sen ³	Krupa et al. 1998
green ash	<i>Fraxinus pennsylvanica</i>	Sen	Krupa and Manning 1988
black huckleberry	<i>Gaylussacia baccata</i>	Sen ²	USDI 2003
witch-hazel	<i>Hamamelis virginiana</i>	Unk	USDI 2003

Continued

Table 7.—continued

Eastern species		Sensitivity	Citation
American holly	<i>Ilex opaca</i>	InSen ¹	Smith 1981
black walnut	<i>Juglans nigra</i>	Unk	
eastern redcedar	<i>Juniperus virginiana</i>	Unk	
tamarack (native)	<i>Larix laricina</i>	Unk	
sweetgum	<i>Liquidambar styraciflua</i>	Sen	Krupa et al. 1998
spicebush	<i>Lindera benzoin</i>	Unk	USDI 2003
yellow-poplar	<i>Liriodendron tulipifera</i>	Sen	Krupa and Manning 1988
maleberry	<i>Lyonia ligustrina</i>	Sen ²	USDI 2003
cucumbertree	<i>Magnolia acuminata</i>	Unk	
apple sp.	<i>Malus sp.</i>	Unk	
blackgum	<i>Nyssa sylvatica</i>	ModSen	Renfro 1987-1992
sourwood	<i>Oxydendrum arboreum</i>	ModSen	Renfro 1987-1992
Virginia creeper	<i>Parthenocissus quinquefolia</i>	Sen ²	USDI 2003
sweet mock orange	<i>Philadelphus coronarius</i>	Sen ²	USDI 2003
Norway spruce	<i>Picea abies</i>	InSen ¹	Smith 1981
white spruce	<i>Picea glauca</i>	InSen ¹	Smith 1981
black spruce	<i>Picea mariana</i>	Unk	
red spruce	<i>Picea rubens</i>	InSen	Eckert et al. 1999
Jack pine	<i>Pinus banksiana</i>	Sen ²	USDI 2003
shortleaf pine	<i>Pinus echinata</i>	ModSen ¹	Smith 1981
table mountain pine	<i>Pinus pungens</i>	Sen	Renfro 1987-1992
red pine	<i>Pinus resinosa</i>	InSen ¹	Smith 1981
pitch pine	<i>Pinus rigida</i>	InSen, Sen ²	Eckert et al. 1999, USDI 2003
eastern white pine	<i>Pinus strobus</i>	Sen	Krupa and Manning 1988
Scotch pine	<i>Pinus sylvestris</i>	ModSen ¹	Smith 1981
loblolly pine	<i>Pinus taeda</i>	Sen	Taylor 1994
Virginia pine	<i>Pinus virginiana</i>	ModSen, Sen ²	Renfro 1987-1992, USDI 2003
American sycamore	<i>Platanus occidentalis</i>	Sen	Krupa and Manning 1988
balsam poplar	<i>Populus balsamifera</i>	Sen ³	Krupa et al. 1998
eastern cottonwood	<i>Populus deltoides</i>	Sen ³	Krupa et al. 1998
bigtooth aspen	<i>Populus grandidentata</i>	Sen ³	Krupa et al. 1998
quaking aspen	<i>Populus tremuloides</i>	Sen	Krupa and Manning 1988
wild plum	<i>Prunus americana</i>	Unk	USDI 2003
pin cherry	<i>Prunus pensylvanica</i>	ModSen	Renfro 1987-1992
black cherry	<i>Prunus serotina</i>	Sen	Krupa and Manning 1988
choke cherry	<i>Prunus virginiana</i>	ModSen	Renfro 1987-1992
white oak	<i>Quercus alba</i>	InSen	Renfro 1987-1992
scarlet oak	<i>Quercus coccinea</i>	ModSen ¹	Smith 1981
northern pin oak	<i>Quercus ellipsoidalis</i>	ModSen ¹	Smith 1981
southern red oak	<i>Quercus falcata</i>	Unk	
shingle oak	<i>Quercus imbricaria</i>	InSen ¹	Smith 1981
bur oak	<i>Quercus macrocarpa</i>	InSen ¹	Smith 1981
pin oak	<i>Quercus palustris</i>	ModSen ¹	Smith 1981
willow oak	<i>Quercus phellos</i>	Unk	
chestnut oak	<i>Quercus prinus</i>	Unk	
northern red oak	<i>Quercus rubra</i>	InSen	Eckert et al. 1999
post oak	<i>Quercus stellata</i>	Unk	
black oak	<i>Quercus velutina</i>	ModSen ¹	Smith 1981
winged sumac	<i>Rhus copallina</i>	Sen ²	USDI 2003
black locust	<i>Robinia pseudoacacia</i>	ModSen, Sen ²	Renfro 1987-1992, USDI 2003
Allegheny blackberry	<i>Rubus allegheniensis</i>	Sen ²	USDI 2003
thornless blackberry	<i>Rubus canadensis</i>	Sen ²	USDI 2003

Continued

Table 7.—continued

Eastern species		Sensitivity	Citation
sand blackberry	<i>Rubus cuneifolius</i>	Sen ²	USDI 2003
black willow	<i>Salix nigra</i>	Unk	
American elder	<i>Sambucus canadensis</i>	Sen ²	USDI 2003
sassafras	<i>Sassafras albidum</i>	Sen	Krupa et al. 1998
common snowberry	<i>Symphoricarpos albus</i>	Sen ²	USDI 2003
northern white-cedar	<i>Thuja occidentalis</i>	InSen	Eckert et al. 1999
American basswood	<i>Tilia americana</i>	InSen ¹	Smith 1981
Chinese tallow	<i>Triadica sebifera</i>	Sen ²	USDI 2003
eastern hemlock	<i>Tsuga canadensis</i>	InSen	Renfro 1987-1992
American elm	<i>Ulmus americana</i>	Unk	
slippery elm	<i>Ulmus rubra</i>	Unk	
northern fox grape	<i>Vitis labrusca</i>	Sen ²	USDI 2003

¹Based on relative sensitivity to acute ozone exposure.

²Based on sensitivity to ambient ozone concentrations in the field and exposure chamber.

³Based on relative sensitivity of genus, not species.

Table 8.—List of western tree and shrub species and their ozone sensitivity

Western species		Sensitivity	Citation
red alder	<i>Alnus rubra</i>	Sen ³	Brace et al. 1996
Sitka alder	<i>Alnus sinuata</i>	Sen	Brace et al. 1996
western serviceberry	<i>Amelanchier alnifolia</i>	ModSen ³	Brace et al. 1996
single-leaf ash	<i>Fraxinus anomala</i>	Sen ⁴	USDI 2003
twinberry	<i>Lonicera involucrata</i>	Sen ⁴	USDI 2003
lodgepole pine	<i>Pinus contorta</i> ¹	ModSen ³	Brace et al. 1996
Jeffrey pine	<i>Pinus jeffreyi</i>	Sen	Miller et al. 1996
ponderosa pine	<i>Pinus ponderosa</i> ²	Sen	Smith 1981
Monterey pine	<i>Pinus radiata</i>	Sen ⁴	USDI 2003
Pacific ninebark	<i>Physocarpus capitatus</i>	Sen ³	Brace et al. 1996
mallow ninebark	<i>Physocarpus malvaceus</i>	Sen ³	Brace et al. 1996
Fremont cottonwood	<i>Populus fremontii</i>	Sen ⁴	USDI 2003
quaking aspen	<i>Populus tremuloides</i>	Sen	Smith 1981
black cottonwood	<i>Populus trichocarpa</i>	ModSen ³	Brace et al. 1996
Douglas-fir	<i>Pseudotsuga menziesii</i>	ModSen ³	Brace et al. 1996
California black oak	<i>Quercus kelloggii</i>	ModSen	Miller et al. 1996
skunk bush	<i>Rhus trilobata</i>	Sen	Temple 2000
thimbleberry	<i>Rubus parviflorus</i>	Sen ⁴	USDI 2003
Gooding's willow	<i>Salix gooddingii</i>	Sen ⁴	USDI 2003
Scouler's willow	<i>Salix scouleriana</i>	Sen ⁴	Brace et al. 1996
willow sp.	<i>Salix sp.</i>	ModSen ⁵	Krupa and Manning 1988
blue elderberry	<i>Sambucus mexicana</i>	Sen	Temple 2000
red elderberry	<i>Sambucus racemosa</i>	ModSen ³	Brace et al. 1996
common snowberry	<i>Symphoricarpos albus</i>	Sen ⁴	USDI 2003
snowberry sp.	<i>Symphoricarpos sp.</i>	Sen ⁵	Smith 1981
western hemlock	<i>Tsuga heterophylla</i>	ModSen ³	Brace et al. 1996
huckleberry	<i>Vaccinium membranaceum</i>	Sen ⁴	USDI 2003
huckleberry sp.	<i>Vaccinium sp.</i>	ModSen ³	Brace et al. 1996

¹*Pinus contorta* var. *latifolia*.

²*Pinus ponderosa* var. *ponderosa*.

³Based on relative sensitivity to acute ozone exposure.

⁴Based on sensitivity to ambient ozone concentrations in the field and exposure chamber.

⁵Based on relative sensitivity of genus, not species.

Appendix III: Documents for Ozone Information Management

Documents to assist FIA regional analysts with ozone information management:

1. Flow of Ozone Data from the Field to the FIA Information Management System
2. Ozone Standard Summary Tables in the FIA Data Base (FIADB)
3. Formulation of the Biosite Index
4. SAS Code for Biosite Tables and Maps
5. Contact List for Ozone Data Management

To obtain the following documents, email Gretchen Smith at gsmith@nrc.umass.edu

1. Computation specifications for derived ozone data in FHM
2. Ozone bioindicator attribute definitions for FIADB
3. Ozone data collection start dates by state and year
4. Crosswalk tables for tracking changes to the ozone sample from 1994 to the present
5. Sample biosite field map
6. National ozone risk map for the sampling period 1994-1998
7. National ozone risk map for the sampling period 1999-2002
8. National map of the 7-year average ozone exposure 1996-2002

Flow of Ozone Data from the Field to FIA Information Management

The goals of ozone information management are to clean up the ozone data files collected by the field crews, correct the regular crew and QA crew data files so they are compatible with the leaf voucher data, and generate ozone summary statistics suitable for further analysis and reporting. The summary statistics are used to generate an ozone risk map and population metrics as described in the main body of this document.

Step 1:

Each Regional Analyst works with the raw data file entered by the field crew and the validation file created by the National Indicator Advisor.

Step 2:

The Regional Analyst/P3 Data Processor in each region takes the raw data files entered by the field crew and the validation file created by the National Indicator Advisor and loads both into NIMS (National Information Management System). The P3 LAB system-checker program determines errors between the validation file and the raw data. Differences between these two files must be resolved at the regional level through direct communication between the National Indicator Advisor and the Regional Analyst. Error resolution requires changes to both the raw data file and the validation file.

Note: It is sometimes helpful to resolve differences between these two files before loading the data into NIMS. Software to assist with this process is available. Once the data are loaded, the checker program is used as a final edit.

Step 3:

The Regional Analyst/P3 Data Processor in each region runs the P3 LAB system-report program on the validated ozone data. The report program creates three ozone standard summary tables: OZONE_PLOT_SUMMARY, OZONE_SPECIES_SUMMARY, and OZONE_BIOSITE_SUMMARY.

Step 4:

The Regional Analyst/P3 Data Processor contacts Brian Cordova, FIA-IM, at: cordovab@unlv.nevada.edu. Each region's data are captured and placed on the national NIMS Web site. Sensitive information is stripped (NULLED), and the remaining information is posted on the national FIA database (FIADB) P3 Web site and the FIADB Data Mart, which is the data distribution system to the public.

Note: Step 2 instructs the Regional Analyst to load the data into NIMS. Until the new TALLY program is completed, the raw TALLY files are parsed using TALLY Cracker and inserted into the LOAD tables in NIMS. The data are moved from the LOAD tables to the NIMS tables through the front-end, which is a graphical interface used to load and drop data, run computations and reports, etc. The front-end is again used to load the validation file and create a report of any errors. In the future, Step 4 will be a direct upload to the NIMS FIA-P3 Web site via the front-end interface.

Additional Steps:

The NIMS ozone summary tables provide biosite summary statistics suitable for preliminary reports at the State and regional levels (see Output Tables and Maps). SAS routines are available that generate additional summary statistics from the validated ozone files. For example, one routine generates all the necessary values to create a summary table that presents numbers of biosites evaluated, number of plants sampled by species, and percentage of sampled plants in each injury severity category. Another SAS routine is available that generates a biosite list with presence or absence of ozone injury to use for an ozone site distribution map. This map is useful for tracking changes over time in the number and distribution of plus ozone sites across a State or region.

The OZONE_BIOSITE_SUMMARY table includes the biosite-level ozone injury index referred to as the Biosite Index (BI). Using a 5-year rolling average of the BI, an FIA Spatial Data Analyst creates the national risk map of probable ozone injury. A new map is produced every year. This map surface is stored in the FIADB Data Mart so that it can be extracted by Regional Analysts, in whole or in part, as needed. FIA Spatial Data Services uses the national map to generate an estimated BI value for every P2 ground plot. This biosite attribute is added to the larger P2 table of plot attributes in the FIADB Data Mart. This will allow FIA analysts to examine relationships between bioindicator attributes and other indicators of tree growth, forest health, and condition. FIA Spatial Data Services also maintains a master list of geographical coordinates for the ozone sampling grid and crosswalk tables that link biosites on the FHM-P3 grid (1994-2001) to biosites on the FIA Ozone Grid (2002-present).

Analysts responsible for 5-year reports or comprehensive regional reports should refer to the main body of the text of this document for detailed guidance on the analytical techniques used to generate FIA P2 plot-level metrics of the ozone data. The companion user guide for the ozone indicator provides (1) examples of output tables and maps using real data, and (2) additional interpretive guidance on the issues associated with ozone air quality and forest health.

Three Ozone Standard Summary Tables in the FIA Data Base

Ozone Species Summary Table

Statecd	number(2)	not null
countycd	number(2)	not null
p3hex	number(7)	not null
p3plot	number(1)	not null
plot	number(8)	not null
measyear	number(4)	not null
bio_species_cd	number	not null
amount_maximum	number	not null
amount_minimum	number	not null
amount_mean	number	not null
severity_maximum	number	not null
severity_minimum	number	not null
severity_mean	number	not null
plants_inj_cnt	number	not null
plants_eval_cnt	number	not null
plants_ratio	number	not null
bio_species_sum	number	not null
bio_species_index	number	not null
elev	number	not null
pltsize	number	not null
aspect	number	not null
terrpos	number	not null
soildpth	number	not null
soildrn	number	not null
soildrn	number	not null
plotwet	number	not null
pltdstrb	number	not null

Ozone Plot Summary Table

Statecd	number(2)	not null
countycd	number(2)	not null
p3hex	number(7)	not null
p3plot	number(1)	not null
plot	number(8)	not null
measyear	number(4)	not null
species_eval_cnt	number	not null
biosite_index	number	not null
elev	number	not null
pltsize	number	not null
aspect	number	not null
terrpos	number	not null
soildpth	number	not null
soildrn	number	not null
soildrn	number	not null
plotwet	number	not null
pltdstrb	number	not null

Ozone Biosite Summary Table

Statecd	number(2)	not null
countycd	number(2)	not null
p3hex	number(7)	not null
location_cnt	number(1)	not null
plot	number(8)	not null
measyear	number(4)	not null
plant_inj_cnt	number	not null
plant_eval_cnt	number	not null
plant_ratio	number	not null
species_eval_cnt	number	not null
biosite_index***	number	not null
svrty_class_zero	number	not null
svrty_class_one	number	not null
svrty_class_two	number	not null
svrty_class_three	number	not null
svrty_class_four	number	not null
svrty_class_five	number	not null

***See notes on the formulation of the biosite_index on the following page.

Formulation of the Biosite Index (BI)

Note: The ozone indicator site-level biosite index was formulated with the assistance of David Randall, Statistician for the USDA Forest Service, Northeastern Area, Washington Office.

Notes on the formulation:

There are 3 components to the formulation: (1) the amount of injury, (2) the severity of injury, and (3) the incidence of injury on the site. The formulation selected associates these three components at the individual plant level. This suggests that the ozone injury response of each individual plant is important. This is biological reality and better than lumping all species together.

The calculation is intuitive. A mean value is calculated that truly represents a proportion of the population at both the plant level and the species level. An arithmetic mean is then taken for the “n” species on the plot.

Notes on method:

Each plant observed by the field crew is rated for the percent of the plant that is injured (i.e., injury amount) and the average severity of injury (i.e., injury severity) using a modified Horsfall-Barrett scale with breakpoints at 6, 25, 50, 75, and 100 percent. This information is used to calculate an injury value for each plant, a mean value for each species, and an overall site mean. The incidence of injury on the site is also considered. The formulation is based on the fact that each individual plant has a unique response to ozone that is dependent on the genotype and microhabitat at the time of exposure.

For each plant:

AMT = injury amount

SEV = injury severity

For each species:

N_1 = the number of injured plants

N_2 = the number of evaluated plants

$A = N_1 / N_2$

$B = \sum[(AMT) (SEV)] / N_1$

bio_species_index = (A)(B)

For each biosite (hexagon number):

N_3 = the number of evaluated species

biosite_index = $\sum(\text{bio_species_index}) / N_3$

Notes on transforming crew values from the ordinal scale to the percent scale:

In the field, the crews estimate the percent injury to the plant and then assign ordinal values that reflect 5 broad classes of injury as follows: 0 = no injury; 1 = 1-6% injury; 2 = 7-25% injury; 3 = 26-50% injury; 4 = 51-75% injury; and 5 = $\geq 75\%$ injury.

In the office, the ordinal codes recorded by the field crews are converted to percentage values representing the midpoint of each injury class as follows: 0 = 0; 1 = 3.5%; 2 = 16%; 3 = 38%; 4 = 63%; and 5 = 88%. Theoretically, the site-value has a range from zero to 100. In reality, the highest values are less than 25 and most are less than 5. A site with no injury has an index of zero.

The measurement intervals on the two scales are different. The intervals on the ordinal scale are equal; those on the percent scale are not. Midpoint values are used rather than ordinal values for the site-level index because (1) the midpoint percentage values bring the reader back to the original scale used by the field crew to rate the injured plants; (2) the percentage values have some intuitive biological relevance, unlike the ordinal scale, which was developed largely as a matter of convenience for the field crews; and (3) readers relate more easily to the percentage values than the ordinal scale.

It is understood that the data transformation introduces some error or misrepresentation into the reporting of the biosite_index. Nevertheless, as long as the reader knows how the biosite-value was calculated, the midpoint percentage is still preferable because it provides a more meaningful image of ozone injury than would be provided by the ordinal scale.

SAS CODE: Biosite Tables

The following routine will generate summary statistics similar to those presented in Table 5.

- The number of plots evaluated by State
- The number of plants sampled by State
- The percent of sampled plants in each HB injury severity class by State

This is a highly defensible presentation of the ozone indicator data for State and regional reports.

```
**** filename o3_summary.sas Jan 2004 Barbara O'Connell****/  
****Program to produce percent injured by severity by State *****/  
*** must first run vouch03.sas, biovch03.sas, and biocor03.sas to create corrected data set****/
```

```
options ls=95 ps=1000 obs=max;
```

```
libname perm 'c:\_barbo\2003\analysis\ozone\';
```

```
data o3sum03; set perm.biocor03; *biocor03 is raw tally files corrected as per validation data;  
if qa_stat=1; *include only the regular crew data;  
if severity =. then severity=0;  
if amount>0 then nplants=1;
```

```
proc sort data=o3sum03; by State hex_num;  
proc freq data=o3sum03;  
tables severity/norow nocol nofreq nocum;  
by State;  
weight nplants;  
title1 '2003 Ozone Severity';
```

```
proc summary data=o3sum03; * to get the number of plots surveyed per State;  
class hex_num;  
id State;  
output out=all03;  
data all03; set all03; total=_freq_; drop _type_ _freq_;  
proc sort data=all03;  
by State hex_num;  
proc summary data=all03;  
class State;  
var hex_num;  
output out=noplots n=;  
proc print;  
title1 '2003 Ozone plots';
```

```
proc summary data=o3sum03; * to get the number of plants sampled;  
class State;  
var nplants;  
output out=spcnt sum=;  
proc print;  
title1 '2003 Nbr. of Plants Sampled';  
run;
```

Note: The SAS routine was written by Barbara M. O'Connell, Forester, USDA Forest Service, Northern Research Station, FIA, Newtown Square, PA.

SAS CODE: Maps

The following routine will generate summary statistics that can be used to create a State or regional map of ozone biomonitoring sites with and without ozone injury. This type of map is useful for documenting status and change in the number and distribution of biosites with ozone injury across a State or region.

```
/*** filename biomap03.sas March 1998 ***/
/** updated Jan 2004 Barbara O'Connell***/
/** must first run vouch03.sas, biovch03.sas, and biocor03.sas to create corrected data set***/

options ls=95 ps=1000 obs=max;

libname perm 'c:\_barbo\2003\analysis\ozone\';

data bio03; set perm.biocor03;
  if qa_stat=1; *include only the regular crew data;

proc summary nway data=bio03; *to create hex_num data set;
  class hex_num;
  id State;
  output out=all;

data all; set all; drop _type_ _freq_;

proc sort data=all; by hex_num ;

data temp; set bio03; if amount > .5; *include only injured data;

proc summary nway data=temp; *to create data set with hex avg injury and total nbr damaged;
  var amount severity;
  class hex_num ;
  output out=damaged mean= ;

data damaged; set damaged; damaged=_freq_; drop _type_ _freq_;
proc sort data=damaged; by hex_num;
data summary;
  merge all damaged;
  by hex_num ;
if damaged=. then damaged=0; if amount < .5 then amount=0;
if severity < .5 then severity=0;
data final; set summary;
if amount = 0 then infect=2;
if amount > .5 then infect=1;

proc sort;
by State hex_num;

proc print r;

title1 'FHM 2003 Bioindicator';
title2 'no. of plants with injury and avg amount/severity per plot';
title4 'injury: 1=injury detected 2=no injury';
var State hex_num infect damaged amount severity;

run;
```

Note: The SAS routine was written by Barbara M. O'Connell, Forester, USDA Forest Service, Northern Research Station, FIA, Newtown Square, PA.

Contact List for Ozone Data Management

National Indicator Advisors:

John Coulston, FS – Southern Research Station, Knoxville, TN, jcoulston@fs.fed.us
Gretchen Smith, UMass-Amherst, gcsmith@nrc.umass.edu

Information Management: For access to validated data files, summary statistics, and map products.

Brian Cordova, UNLV, cordovab@unlv.nevada.edu
Chuck Liff, FS – Rocky Mountain Research Station, Ogden, UT, cliff@fs.fed.us
John Coulston, FS – Southern Research Station, Knoxville, TN, jcoulston@fs.fed.us

National QA: For information on QA analyses of the field data at the national level.

Jim Pollard, UNLV, pollardj@unlv.nevada.edu
Bill Smith, FS – RTP, bdsmith@fs.fed.us

Regional Analysts: For information on analyses and reports and for access to regional files.

Barb O'Connell, FS – Northern Research Station, Newtown Square, PA, boconnell@fs.fed.us
Randy Morin, FS - Northern Research Station, Newtown Square, PA, rsmorin@fs.fed.us
Ed Jepsen, WI Department of Natural Resources, jepsee@dnr.state.wi.us
Mark Hansen, FS – Northern Research Station, St. Paul, MN, mhansen01@fs.fed.us
John Simpson, FS – Southern Research Station, Knoxville, TN, jsimpson03@fs.fed.us
Anita Rose, FS – Southern Research Station, Knoxville, TN, anitarose@fs.fed.us
Mark Ruby, FS – Rocky Mountain Research Station, Ogden, UT, mruby@fs.fed.us
James Menlove, FS - Rocky Mountain Research Station, Ogden, UT, jmenlove@fs.fed.us
Ron Wanek, FS – Pacific Northwest Research Station, Portland, OR, rwane@fs.fed.us
Sally Campbell, FS – Pacific Northwest Research Station, Portland, OR, Scampbell01@fs.fed.us

Regional Advisors: For information on ozone implementation, training, and QA procedures.

Gretchen Smith, UMass-Amherst, gcsmith@nrc.umass.edu
Ed Jepsen, WI DNR, jepsee@dnr.state.wi.us
Dan Stratton, FS – Southern Research Station, Knoxville, TN, dstratton@fs.fed.us
Bill Dunning, FS – Rocky Mountain Research Station, Ogden, UT, bdunning@fs.fed.us
Sally Campbell, FS – Pacific Northwest Research Station, Portland, OR, scampbell01@fs.fed.us

Where to find more information on the biomonitoring program and ozone data.

<http://nrs.fs.fed.us/fia/topics/ozone/default.asp> This is the ozone indicator home page. Click on links to data, methods, ozone sensitive species, publications, contacts, and other information.

<http://fia.fs.fed.us/> This is the FIA home page. Enter 'ozone' in the search window for access to the ozone indicator home page; Click on FIA Data and Tools for access to field data, standard summary tables, and the FIADB user guide for field attributes. Click on FIA Library for access to the field manual and fact sheet.

<http://fhm.fs.fed.us/> This is the FHM home page. It includes fact sheets for all of the P3 forest health indicators.

FIA uses the Oracle Database Management System to process and store the ozone indicator data (also called NIMS – National Information Management System). Raw data from field-collected tally files and sample voucher validation files are loaded onto five standard tables (OZONE_PLOT_TBL, OZONE_VISIT, OZONE_SPECIES, OZONE_PLOT_NOTES, and OZONE_VALIDATION). Further processing computes indices and creates three standard summary tables (OZONE_BIOSITE_SUMMARY, OZONE_PLOT_SUMMARY, and OZONE_SPECIES_SUMMARY), which are used as the presentation data in the FIADB (FIA database) after sensitive information has been stripped. The database structure is such that flat files can be easily produced for users who do not have access to, or the capability of, database management on their own computers. ASCII data files (1994 to present), two core ozone maps, and a core list of ozone-sensitive tree and shrub species are available

for download from the FIADB Data Mart (the data distribution system to the public). The first map product is the national ozone risk map that provides an interpolated surface of probable ozone injury across the landscape. The second map product is an interpolated surface of ambient ozone concentrations. Data users select their area of interest (e.g., state, region, or eco-region) from these two map products, and use the procedures outlined in the ozone estimation document to calculate and interpret population metrics for the ozone indicator. If you have trouble accessing Web sites or data files, contact the National Ozone Advisor, the FIA analyst in your region, or Brian Cordova at cordovab@unlv.nevada.edu.

Smith, Gretchen C.; Smith, William D.; Coulston, John W. 2007. **Ozone Bioindicator Sampling and Estimation**. Gen. Tech. Rep. NRS-20. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 34 p.

Ozone is an important forest stressor that has been measured at known phytotoxic levels at forest locations across the United States. The percent forest exhibiting negative impacts from ozone air pollution is one of the Montreal Process indicators of forest health and vitality. The ozone bioindicator data of the U.S. Forest Service Forest Inventory and Analysis Program (FIA) are the only source of information available that documents plant injury from air pollution using consistent protocols. This document introduces the FIA ozone indicator and describes the sampling and estimation procedures of the national biomonitoring program. We provide background material on ozone, examples of bioindicator summary statistics, a description of spatial interpolation, and methods to estimate status and change in forested areas with respect to the occurrence of ozone injury from ambient ozone concentrations. The goal is to provide guidance to analysts and researchers on ways to incorporate ozone bioindicator data into reports and research studies. Periodic recommendations to analysts on improved analytical techniques will be made. Analysts are encouraged to consult the companion user guide for additional guidance on interpreting the ozone biomonitoring data and reporting on the issue of ozone and forest health for the FIA program.

KEY WORDS: air quality, bioindicator species, biomonitoring, forest health, kriging, ozone, ozone sensitive, risk assessment, spatial interpolation

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternate means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202)720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, DC 20250-9410, or call (800)795-3272 (voice) or (202)720-6382 (TDD). USDA is an equal opportunity provider and employer.



Printed on Recycled Paper



Capitalizing on the strengths of existing science capacity in the Northeast and Midwest to attain a more integrated, cohesive, landscape-scale research program