


Pasture Measurement and Perennials




Measuring Pasture Biomass



Maturity Map
Please hover over the blo
scale




OV Farm Pasture Report



Est Dry Matter/Paddock

Pre-grazing target




Target residual







Date: 8/9/2022




Est Dry Matter/Paddock








Date: 10/28/2022



Est Dry Matter/Paddock












7/5/2021




Est Dry Matter/Paddock



Date: 8/9/2022



Est Dry Matter/Paddock



World Forage Analysis Superbowl

2022 Forage Analysis Superbowl

FINALIST NAME
Curvin Brubacher
W12074 County N
Stanley WI 54768

DRY MATTER.....	76.48 %
CRUDE PROTEIN.....	17.16 %
NEUTRAL DETERGENT FIBER.....	41.81 %
ASH.....	10.90 %
30HR NDFD % of NDF (in vitro).....	57.49
RELATIVE FORAGE QUALITY (RFQ).....	165
MILK 2006 TDN	63.68
MILK 2006 MILK PER TON.....	3219
VARIETY.....	L310
FINAL SCORE	75.3
VISUAL SCORE.....	63.7


CLASS G1-GRASS HAY PLACING: 9







Est Dry Matter/Paddock

















Est Dry Matter/Paddock









7/5/2021




Est Dry Matter/Paddock






OV Farm Pasture Report



Est Dry Matter/Paddock

Pre-grazing target



Target residual









Figure 3.2. The time-independent relationship between instantaneous herbage accumulation rate (HAR_i) (from Fig. 3.1b) and herbage mass above ground-level (from Fig. 3.1a). The maximum instantaneous herbage accumulation rate (HAR_{i-max}) was 33.1 kg DM ha⁻¹ d⁻¹ and the critical range of herbage mass for >90% of maximum instantaneous herbage accumulation rate ($HAR_{i-90\%}$) (29.8 kg DM ha⁻¹ d⁻¹) was between 2760 and 4170 kg DM ha⁻¹.



Date: 8/9/2022



Est Dry Matter/Paddock

