



## **INTERPRETING THE RESULTS OF SOIL TESTS FOR HEAVY METALS**

by Vern Grubinger, Don Ross, and Joshua Faulkner

Agricultural soils normally contain low background levels of heavy metals. Contamination from industrial activities or byproducts can increase the natural levels of heavy metals in soil, creating a health hazard to people, livestock and plants. Fertilizers and other soil amendments also add small amounts of heavy metals to the soil, which can build up over time with repeated applications.

The actual toxicity of a heavy metal will be affected by soil texture, organic matter, and pH. The health effects of exposure to heavy metals depend on the amount and duration of exposure, i.e. the volume of contaminated soil or food consumed over time.

It is not clear exactly what levels of heavy metals in soil are safe or unsafe, so the following information is provided only to help you understand your test results and the relative level of risk they represent. In soils with elevated heavy metal levels, which may pose higher levels of risk, you should consider whether remedial actions are appropriate, or whether crops should be grown at all.

### **UVM Heavy Metals Soil Test**

The test provided by UVM is only a screen for heavy metals and does not measure the actual total metal content of the soil. This low-cost test uses a weak acid to extract heavy metals. The amount of metal extracted is roughly proportional to the total amount present. Maximum levels for heavy metals in soils established by regulatory agencies are based on total heavy metal content (see below) and require a more involved and expensive test. If your UVM test results indicate an elevated level of heavy metal(s), you should consider submitting another sample for a total heavy metal analysis (see more information specific to lead, below).

### **Interpreting TOTAL Heavy Metals Soil Test Results**

The US Environmental Protection Agency (EPA) and NY Department of Environmental Conservation (NYS DEC) have guidelines for determining the safety of various land uses based on total soil metal concentrations. Table 1 shows these limits, which are used to guide clean-up efforts. EPA levels are used to guide clean-up efforts of contaminated sites; NYS DEC levels are based on removing human health risks; unrestricted use includes agriculture.

| <b>Table 1. Levels of heavy metals in soil used to guide cleanup and land use decisions (mg/kg)</b> |                               |                   |                 |
|---|-------------------------------|-------------------|-----------------|
|   | US EPA                        | NYS DEC           |                 |
|   | Soil level requiring clean-up | Unrestricted use* | Residential use |
| Copper (Cu)   | --                            | 50                | 270             |
| Cadmium (Cd)  | 70                            | 2.5               | 2.5             |
| Chromium (Cr)   | 230                           | 30                | 36              |
| Nickel (Ni)   | 1600                          | 30                | 140             |
| Lead (Pb)   | 400                           | 63                | 400             |
| Zinc (Zn)   | 23,600                        | 109               | 2200            |
| *Includes agricultural use.   |                               |                   |                 |

### **Lead is a Special Concern**

There has been a lot of attention paid to lead levels in soil because it is well-known to cause adverse health effects and is relatively widespread because of its historical use in many commercial products, from gasoline to paint.

Table 2 shows the guidelines for garden soil use based on **total** lead content that have been developed by the states of New Jersey, Pennsylvania, and Vermont. Lead concentrations reported on your UVM Soil Test report reflect extractable lead, not total lead.

To estimate total lead concentration, multiply the extractable lead concentration on the UVM test report by 10. For a more precise determination of total lead, samples can be submitted to the UVM soil testing lab for a total lead analysis.

| <b>Table 2. Soil lead contamination levels and recommended actions.</b> |                           |           |           |  |
|---|---------------------------|-----------|-----------|--|
| Contamination level   | Total Lead in soil, mg/kg |           |           | Recommended Action   |
|   | PA                        | NJ        | VT        |  |
| none / very low   | < 150                     | < 100     | <41*      | No need to be concerned about lead exposure.   |
| low / elevated  | 150 - 400                 | 100 - 300 | 41 – 200  | Conduct best management practices (BMPs) to minimize lead exposure from vegetable gardens: apply phosphate fertilizer, maintain high pH for fruiting vegetables, keep soil mulched to minimize dust and lead inhalation. |
| medium / significant  | 400 - 1000                | 300 - 400 | 200 – 400 | Conduct BMPs; do not grow leafy vegetables.  |
| high / cleanup  | > 1000                    | > 400     | >400      | Do not grow a vegetable garden.<br><br>Contact local health department for lead abatement measures.  |

\*VTDEC has adopted 41 ppm as the residential statewide soil lead background threshold level.  
[https://dec.vermont.gov/sites/dec/files/documents/Vermont.Soil\\_.Full\\_.Report.pdf](https://dec.vermont.gov/sites/dec/files/documents/Vermont.Soil_.Full_.Report.pdf)

**Best Management Practices for Soils with Elevated Levels of Heavy Metals**

Although heavy metals remain in soil for a very long time, there are some steps that can be taken to reduce the level of risk they pose. In some cases, heavy metal concentrations can be ‘diluted’ with deep tillage; for example, to distribute contaminated surface sediment that was deposited by flooding. In garden plots, dilution can be achieved by the addition of uncontaminated soil. Adding organic matter to the soil can help ‘tie up’ heavy metals chemically, reducing their availability for potential plant uptake. Similarly, liming to a neutral pH and maintaining optimal soil phosphorus levels can reduce heavy metal availability to plants. For some heavy metals, such as lead, there is little evidence that it is accumulated within crops; the main health hazard is through soil ingestion and inhalation. Soils that are high in heavy metals pose a greater health risk to children than to adults because children are still growing, and they are more likely to ingest soil directly.

To reduce health risks in soils with elevated heavy metal content, food crops should be thoroughly washed to remove as much soil as possible. The outer leaves of leafy greens should be removed, and root crops should be peeled to further reduce risk.

## **References**

Angima, S. Toxic heavy metals in farm soil. Oregon State University Small Farm News, summer 2010. <https://smallfarms.oregonstate.edu/smallfarms/toxic-heavy-metals-farm-soil>

Hamel, S., J. Heckman, and S. Murphy. 2010. Lead contaminated soil: minimizing health risks. Fact sheet FS336. Rutgers, the State University of New Jersey, New Jersey Agricultural Experiment Station. <http://njaes.rutgers.edu/pubs/publication.asp?pid=FS336>

Hoskins, B. 2022. General Guidelines for Soil Lead Contamination. University of Maine. <https://umaine.edu/soiltestinglab/wp-content/uploads/sites/227/2022/06/lead-guidelines2022.pdf>

NYS DEC. 2006. New York State Brownfield Cleanup Program Development of Soil Cleanup Objectives Technical Support Document. New York State Department of Environmental Conservation and New York State Department of Health, Albany, NY. [https://www.dec.ny.gov/docs/remediation\\_hudson\\_pdf/part375.pdf](https://www.dec.ny.gov/docs/remediation_hudson_pdf/part375.pdf)

Stehouwer, R. and K. Macneal, 1999. Lead in residential soils: sources, testing, and reducing exposure. Pennsylvania State University, College of Agricultural Sciences, Cooperative Extension. <https://extension.psu.edu/lead-in-residential-soils-sources-testing-and-reducing-exposure>

Pettinelli, D. Soil Lead Interpretation Sheet. Soil Nutrient Analysis Laboratory, University of Connecticut, Storrs. <https://soiltesting.cahn.uconn.edu/wp-content/uploads/sites/3514/2022/06/Lead-Interpretation-Sheet.pdf>

*The information in this document reflects our best effort to interpret regulatory guidelines and scientific research, and to translate this into practical management options. However, growers are fully responsible for their own management decisions, for the quality of the food they sell, and for compliance with all applicable laws and regulations.*

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. University of Vermont Extension, Burlington, Vermont. University of Vermont Extension, and U.S. Department of Agriculture, cooperating, offer education and employment to everyone without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or familial status.

*Revised May 5, 2023*