

IS PULSAR B0656+14 A VERY NEARBY ROTATING RADIO TRANSIENT?

P. WELTEVREDE,¹ B. W. STAPPERS,^{1,2} J. M. RANKIN,^{3,4} AND G. A. E. WRIGHT^{4,5}

Received 2006 April 26; accepted 2006 June 1; published 2006 July 7

ABSTRACT

The recently discovered RRAT sources are characterized by very bright radio bursts that, while being periodically related, occur infrequently. We find bursts with the same characteristics for the known pulsar B0656+14. These bursts represent pulses from the bright end of an extended smooth pulse-energy distribution and are shown to be unlike giant pulses, giant micropulses, or the pulses of normal pulsars. The extreme peak fluxes of the brightest of these pulses indicate that PSR B0656+14, were it not so near, could only have been discovered as an RRAT source. Longer observations of the RRATs may reveal that they, like PSR B0656+14, emit weaker emission in addition to the bursts.

Subject headings: pulsars: general — radiation mechanisms: nonthermal — stars: individual (PSR B0656+14)

Online material: color figure

1. INTRODUCTION

PSR B0656+14 is one of three nearby pulsars in the middle-range in which pulsed high-energy emission has been detected. These are commonly known as “The Three Musketeers” (Becker & Trümper 1997), the other two being Geminga and PSR B1055–52. PSR B0656+14 was included in a recent extensive survey of subpulse modulation in pulsars in the northern sky at the Westerbork Synthesis Radio Telescope by Weltevrede et al. (2006). In the single pulses analyzed for this purpose, the unusual nature of this pulsar’s emission was very evident, especially the brief yet exceptionally powerful bursts of radio emission.

These extreme bursts of radio emission of PSR B0656+14 are similar to those detected in the recently discovered population of bursting neutron stars. These rotating radio transients (RRATs; McLaughlin et al. 2006) typically emit detectable radio emission for less than one second per day, causing standard periodicity searches to fail in detecting the rotation period. From the greatest common divisor of the time between bursts, a period has been found for 10 of the 11 sources. The periods (between 0.4 and 7 s) suggest these sources may be related to the radio-quiet X-ray populations of neutron stars, such as magnetars (Woods & Thompson 2006) and isolated neutron stars (Haberl 2004). However, Popov et al. (2006) have shown that the estimated formation rate of magnetars is too low. Furthermore, the spectrum of the only RRAT for which an X-ray counterpart has so far been detected (Reynolds et al. 2006) seems to be too cool, too thermal, and too dim for a magnetar, but is consistent with a cooling middle-aged neutron star such as PSR B0656+14 (Shibanov et al. 2006). Also, the pulse period and the slowdown rate of PSR B0656+14, as well as the derived surface magnetic field strength and characteristic age, are within the range of measured values for RRATs.

2. OBSERVATIONS

The results in this paper are based on an archival and a new observation made using the 305 m Arecibo telescope on 2003 July 20 and 2005 April 30, respectively. Both observations had a center frequency of 327 MHz and a bandwidth of 25 MHz. Almost 25,000 and 17,000 pulses with a sampling time of 0.5125 and 0.650 ms were recorded using the Wideband Arecibo Pulsar Processor (WAPP⁶) for the 2003 and 2005 observations, respectively. The Stokes parameters have been corrected off-line for dispersion, Faraday rotation, and various instrumental polarization effects.

The data were in some instances affected by radio frequency interference, but this could relatively easily be removed by excluding the pulses with the highest rms of the off-pulse noise (about 1% of the data in both observations) from further analysis. The results derived from both observations (of which the one from 2005 is relatively clean) are very similar, making us confident in the results.

The observations are not flux calibrated, but are sufficiently long to get a pulse profile with high precision. From its shape it follows that the peak flux of the profile is about 17 times that of the integrated flux density over the entire pulse phase. The average flux at our observing frequency is estimated to be 7.2 mJy, based on the measurement of the spectral index (–0.5) and flux density by Lorimer et al. (1995) at 408 MHz. Therefore the peak flux of the profile is approximately 0.12 Jy. The scintillation bandwidth of PSR B0656+14 is much smaller than the observing bandwidth, so no intensity fluctuations appear in the data as a function of time due to interstellar scintillation.

3. THE RADIO BURSTS OF PSR B0656+14

To characterize the bright pulses of PSR B0656+14, the pulse-energy distribution is calculated (see Fig. 1). In this plot the energies are normalized to the average pulse energy $\langle E \rangle$. The brightest measured pulse is $116\langle E \rangle$, which is exceptional for regular radio pulsars. Based on the energy of these pulses alone, PSR B0656+14 would fit into the class of pulsars that emit so-called giant pulses (e.g., Cairns 2004). About 0.3% of the pulses of PSR B0656+14 are brighter than $10\langle E \rangle$, which is the working definition of giant pulses. Nevertheless, there are important differences between giant pulses and the bright

¹ Astronomical Institute “Anton Pannekoek,” University of Amsterdam, Kruislaan 403, 1098 SJ Amsterdam, Netherlands; wltvrede@science.uva.nl.

² Stichting ASTRON, Postbus 2, 7990 AA Dwingeloo, Netherlands; stappers@astron.nl.

³ Physics Department, 405 Cook Physical Science Building, University of Vermont, Burlington, VT 05405; joanna.rankin@uvm.edu.

⁴ Visiting Astronomer, Astronomical Institute “Anton Pannekoek,” University of Amsterdam.

⁵ Astronomy Centre, University of Sussex, Falmer BN1 9QJ, UK; g.wright@sussex.ac.uk.

⁶ See <http://www.naic.edu/~wapp>.




FIG. 1.—Pulse-energy distribution of the 2005 observation of PSR B0656+14 (*solid line*) and the off-pulse distribution (*dashed line*). The brightest pulse is about 116 times stronger than the average, which is outside the plotted energy range.

bursts of PSR B0656+14. The bursts of PSR B0656+14 have timescales that are much longer than the nanosecond timescale observed for giant pulses (e.g., Soglasnov et al. 2004; Hankins et al. 2003), do not show a power-law energy-distribution, are not confined to a narrow pulse window, and are not associated with an X-ray component. This suggests differing emission mechanisms for the classical giant pulses and the bursts of PSR B0656+14. Also, the possible correlation between emission of giant pulses and high magnetic field strengths at the light cylinder (around 10^5 G; Cognard et al. 1996) clearly fails for PSR B0656+14 (766 G). However, giant pulses have been claimed in other (slow) pulsars that also easily fail this test (e.g., Kuzmin & Ershov 2004; Kramer et al. 2003), and for millisecond pulsars a high magnetic field strength at the light cylinder seems to be a poor indicator of the rate of emission of giant pulses (Knight et al. 2006).

The bursts of PSR B0656+14 are even more extreme when we consider their peak fluxes (see Fig. 2). The highest measured peak flux of a burst is 420 times the average peak flux of the pulsed emission, which is an order of magnitude brighter than the giant micropulses observed for the Vela pulsar (Johnston et al. 2001) and PSR B1706–44 (Johnston & Romani 2002). Giant micropulses are not necessarily extreme in the sense of having a large integrated energy (as required for giant pulses),




FIG. 2.—*Dashed line*: Average profile of our 2005 observation. *Solid line*: Ratio between the peak flux of the brightest burst at each longitude and the average peak flux of the profile.

but their peak-flux densities are very large. Not only are the bursts of PSR B0656+14 much brighter (both in peak flux and integrated energy) than those found for giant micropulses; they are also not confined in pulse longitude and they do not show a power-law energy distribution as the giant pulses and micropulses do.

At the leading edge of the profile we detected a burst with an integrated pulse energy of $12.5\langle E \rangle$. What makes this pulse so special is that it has a peak flux that is 2000 times that of the average emission at that pulse longitude (Fig. 3, *left*). Its dispersion track exactly matches what is expected for this pulsar (Fig. 3, *middle*), proving that this radio burst is produced by the pulsar. Note that the effect of interstellar scintillation is also clearly visible (different frequency channels have different intensities) and that the dispersion track is the same for the two pulses in the center of the profile. This burst demonstrates that the emission mechanism operating in this pulsar is capable of producing intense sporadic bursts of radio emission even at early phases of the profile. There are only two bursts with a peak flux above the noise level detected at the longitude of the peak of this pulse out of the total of almost 25,000 pulses (see Fig. 3, *right*). This implies either that these two bursts belong to an extremely long tail of the distribution, or that there is no emission at this longitude other than such sporadic bursts.




FIG. 3.—Bright radio burst detected at the leading edge of the pulse profile in the 2003 observation. *Left*: The burst (*solid line*) compared with the average pulse profile (*dashed line*). *Middle*: Same burst, but now with frequency resolution. The data in this plot are not de-dispersed, and their dispersion track matches exactly what is expected for the known dispersion measure of this pulsar (*dashed line*). *Right*: Longitude-resolved energy distribution at the longitude of the peak of the strong pulse (*solid line*) and the off-pulse distribution (*dashed line*). The peak fluxes (F_i) are normalized to the average peak flux of the profile ($\langle F_p \rangle$). [See the electronic edition of the *Journal* for a color version of this figure.]

TABLE 1
COMPARISON OF THE PEAK FLUX OF THE BRIGHTEST BURSTS
OF PSR B0656+14 AND THOSE OF THE RRATs

Name	S_{peak} (mJy)	D (kpc)	L_{peak} (Jy kpc ²)
B0656+14	59000	0.288	4.1
J0848-43	100	5.5	3.0
J1317-5759	1100	3.2	11
J1443-60	280	5.5	8.5
J1754-30	160	2.2	0.77
J1819-1458	3600	3.6	47
J1826-14	600	3.3	6.5
J1839-01	100	6.5	4.2
J1846-02	250	5.2	6.8
J1848-12	450	2.4	2.6
J1911+00	250	3.3	2.7
J1913+1333	650	5.7	21

NOTES.—Here S_{peak} is the peak flux of the brightest detected burst for each source, D the distance, and $L_{\text{peak}} = S_{\text{peak}}D^2$ the peak luminosity of the brightest detected burst.

4. THE RRAT CONNECTION

It is unclear how the extreme pulses of PSR B0656+14 fit into the zoo of apparently different emission types of radio pulsars. They are brighter than the giant micropulses and not constrained to a particular pulse longitude, and despite being energetic enough, they are too broad to be characterized as classical giant pulses. However, the observational facts are that PSR B0656+14 occasionally emits extremely bright bursts of radio emission that are short in duration. One cannot help but see the similarities with the RRATs.

One important question is whether the luminosities of the bursts of the relatively nearby PSR B0656+14 (288 pc; Brisken et al. 2003) and those of the RRATs are comparable (there is no indication that the spatial distributions of RRATs and PSRs are different; McLaughlin et al. 2006). The brightest burst we found in the center of the profile has a peak flux that is 420 times the average peak flux. With an average peak flux of 0.12 Jy (see § 2), this corresponds to a peak flux of 50 Jy. If one compares luminosities (Table 1), one can see that the brightest burst of PSR B0656+14 is as luminous as those of 4 of the 11 RRATs, and therefore very typical for these sources.

It is interesting to compare not only the luminosities of the bursts but also their peak-flux distributions. Although the slope

of the top end of the distribution of PSR B0656+14 is in the range of the giant pulses (between -2 and -3), it is better described by a lognormal than by a power-law distribution. This again suggests that the bright bursts of PSR B0656+14 are different from the classical giant pulses. The top end of the RRAT distribution with the highest number of detections seems to be harder (with a slope -1), but for the other RRATs this is as yet unclear. For instance, the tail of the distribution of PSR B0656+14 seems to be consistent with the distribution of the RRAT with the second highest number of detections.

Normal periodicity searches failed to detect the RRATs, which places an upper limit on the average peak-flux density of weak pulses among the detected bursts of about 1 : 200 (McLaughlin et al. 2006). Because the brightest burst of PSR B0656+14 exceeds the underlying peak flux by a much greater factor, PSR B0656+14 could have been identified as an RRAT, were it not so nearby. Were it located 12 times farther away (thus farther than five of the RRATs), we estimate that only one burst per hour would be detectable (the RRATs have burst rates ranging from one burst every 4 minutes to one every 3 hours). The typical burst duration (about 5 ms) of PSR B0656+14 also matches that of the RRATs (between 2 and 30 ms).

Were PSR B0656+14 12 times more distant, the rms of the noise would increase by a factor 144 relative to the strength of the pulsar signal. When we artificially add Gaussian-distributed noise at this level to the observation, we find no sign of the pulsar's (2.6 Hz) rotation frequency in 35 minute segments of the data⁷ (Fig. 4, *left*) yet the brightest pulse (Fig. 4, *right*) is easily detected (18σ). Only in the spectrum of the whole 1.8 hr observation—which is significantly longer than the continuous observations of the Parkes Multibeam Survey—is the periodicity marginally detectable. This means that a distant PSR B0656+14 could only be found as an RRAT in a survey using Arecibo, unless the pointings were unusually long.

5. IMPLICATIONS AND DISCUSSION

We have shown that PSR B0656+14 could have been identified as an RRAT, had it been at the typical distance of the

⁷ For telescopes with a lower sensitivity than Arecibo (e.g., Parkes), then even if PSR B0656+14 were quite a bit closer it would not reveal its periodicity in a similarly long observation.




FIG. 4.—*Left*: Power spectrum of the first 35 minutes of the 2005 observation of PSR B0656+14 with artificially increased noise corresponding to a 12 times larger distance of this pulsar. The dashed lines indicates the rotation frequency of this pulsar. *Right*: Brightest recorded pulse in the same piece of data, again for a 12 times larger distance.

known RRATs. We have no way of telling whether its capacity to produce intense bursts of emission right across its profile is related to its age, period, inclination, or even its immediate Galactic environment, since this behavior has been found in no other pulsar. The pulse-energy distribution is not a power law, but is better fitted by a lognormal distribution and such distributions are thought to be common for pulsars (e.g., Cairns et al. 2004).

In a study of 32 pulsars, Ritchings (1976) found that PSR B0950+08 shows the highest degree of pulse-to-pulse intensity variation. Nevertheless, the brightest pulse found in an extensive study of its field statistics by Cairns et al. (2004) is approximately $5\langle E \rangle$. Vela does not show pulses brighter than $10\langle E \rangle$, and only 0.5% of the pulses are brighter than $3\langle E \rangle$ (Johnston et al. 2001). For PSR B0656+14, 4% of the pulses are brighter than $3\langle E \rangle$. Therefore the emission of PSR B0656+14 appears to be extremely erratic compared with both normal pulsars and pulsars with giant micropulses.

One could wonder whether more known pulsars show a similar kind of sporadic bright emission. To answer this question one should analyze the pulse-energy distribution of a sample of pulsars. A complication would be that longer than typical observations are required to detect the presence of a tail of strong pulses. In a large survey of subpulse modulation by Weltevrede et al. (2006), this pulsar was the only pulsar that showed clear evidence for this kind of sporadic emission.

Our identification of PSR B0656+14 with RRATs implies that at least some RRATs could be sources that emit pulses continuously, but over an extremely wide range of energies. This is in contrast to a picture (predicted by Zhang et al. 2006) of infrequent powerful pulses with otherwise no emission.

Therefore, if it indeed turns out that PSR B0656+14 (despite its relatively short period) is a true prototype for an RRAT, we can expect future studies to demonstrate that RRATs emit much weaker pulses among their occasional bright bursts. We would also predict that their integrated profiles will be found to be far broader than the widths of the individual bursts, and will need many thousands of bursts to stabilize.

Hopefully, radio observations of RRATs will soon be able to test these predictions. These, together with the detection of more RRATs and potentially their high-energy counterparts, will shed light on their true nature. The transient nature of these sources makes them difficult to detect. However, it is likely that the Galactic population exceeds that of the “normal” radio pulsars (McLaughlin et al. 2006). Thus surveys with long pointings, such as those planned with LOFAR, or many observations of the same region of sky are required. Surveys at low frequencies will also be more sensitive to nearer RRATs, as the greater degree of dispersion will allow them to be more easily distinguished from radio frequency interference.

G. A. E. W. and J. M. R. thank the Netherlands Foundation for Scientific Research (NWO) and the Anton Pannekoek Institute, Amsterdam, for their kind hospitality, and G. A. E. W. the University of Sussex for a Visiting Fellowship. Portions of this work (for J. M. R.) were carried out with support from US National Science Foundation grants AST 99-87654 and 00-98685. Arecibo Observatory is operated by Cornell University under contract to the US National Science Foundation. The Westerbork Synthesis Radio Telescope is operated by the AS-TRON (Netherlands Foundation for Research in Astronomy) with support from NWO.

REFERENCES

- Becker, W., & Trümper, J. 1997, *A&A*, 326, 682
 Brisken, W. F., et al. 2003, *ApJ*, 593, L89
 Cairns, I. H. 2004, *ApJ*, 610, 948
 Cairns, I. H., Johnston, S., & Das, P. 2004, *MNRAS*, 353, 270
 Cognard, I., et al. 1996, *ApJ*, 457, L81
 Haberl, F. 2004, *Adv. Space Res.*, 33, 638
 Hankins, T. H., et al. 2003, *Nature*, 422, 141
 Johnston, S., & Romani, R. 2002, *MNRAS*, 332, 109
 Johnston, S., van Straten, W., Kramer, M., & Bailes, M. 2001, *ApJ*, 549, L101
 Knight, H. S., Bailes, M., Manchester, R. N., Ord, S. M., & Jacoby, B. A. 2006, *ApJ*, 640, 941
 Kramer, M., et al. 2003, *A&A*, 407, 655
 Kuzmin, A. D., & Ershov, A. A. 2004, *A&A*, 427, 575
 Lorimer, D. R., et al. 1995, *MNRAS*, 273, 411
 McLaughlin, M. A., et al. 2006, *Nature*, 439, 817
 Popov, S. B., Turolla, R., & Possenti, A. 2006, *MNRAS*, 369, L23
 Reynolds, S. P., et al. 2006, *ApJ*, 639, L71
 Ritchings, R. T. 1976, *MNRAS*, 176, 249
 Shibanov, Y. A., et al. 2006, *A&A*, 448, 313
 Soglasnov, V. A., et al. 2004, *ApJ*, 616, 439
 Weltevrede, P., Edwards, R. T., & Stappers, B. W. 2006, *A&A*, 445, 243
 Woods, P. M., & Thompson, C. 2006, in *Compact Stellar X-Ray Sources*, ed. W. H. G. Lewin & M. van der Klis (Cambridge: Cambridge Univ. Press), in press (astro-ph/0406133)
 Zhang, B., Gil, J., & Dyks, J. 2006, *ApJL*, submitted (astro-ph/0601063)